

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 39/385, 39/39, C07K 1/10	A1	(11) International Publication Number: WO 98/34642
		(43) International Publication Date: 13 August 1998 (13.08.98)
(21) International Application Number: PCT/US98/02194 (22) International Filing Date: 3 February 1998 (03.02.98) (30) Priority Data: 08/796,316 7 February 1997 (07.02.97) US (71) Applicant: FORDHAM UNIVERSITY [US/US]; 441 East Fordham Road, Bronx, NY 10471 (US). (72) Inventor: SRIVASTAVA, Pramod, K.; 4601 Henry Hudson Parkway, Riverdale, NY 10471 (US). (74) Agents: ANTLER, Adriane, M. et al.; Pennie & Edmonds LLP, 1155 Avenue of the Americas, New York, NY 10036 (US).		(81) Designated States: AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, EE, GE, GH, GW, HU, ID, IL, IS, JP, KG, KP, KR, KZ, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UZ, VN, YU, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published With international search report.
(54) Title: ADOPTIVE IMMUNOTHERAPY UTILIZING HEAT SHOCK/STRESS PROTEIN-PEPTIDE COMPLEXES FOR PREVENTION/TREATMENT OF CANCER OR INFECTIOUS DISEASES		
(57) Abstract The present invention relates to methods and compositions for eliciting an immune response and the prevention and treatment of primary and metastatic neoplastic diseases and infectious diseases. The methods of the invention comprise administering a composition comprising an effective amount of a complex, in which the complex consists essentially of a heat shock protein (hsp) noncovalently bound to an antigenic molecule in combination with administering antigen presenting cells sensitized with complexes of hsps noncovalently bound to an antigenic molecule. "Antigenic molecule" as used herein refers to the peptides with which the hsps are endogenously associated <i>in vivo</i> as well as exogenous antigens/immunogens (i.e.), with which the hsps are not complexed <i>in vivo</i> or antigenic/immunogenic fragments and derivatives thereof. In a preferred embodiment, the complex is autologous to the individual. In a specific embodiment, the effective amounts of the complex when administered intradermally are in the range of 0.1 to 9.0 micrograms for complexes comprising hsp70, 5 to 49 micrograms for hsp90, and 0.1 to 9.0 micrograms for gp96. In another embodiment, the effective amounts of the complex when administered subcutaneously are in the range of 10 to 600 micrograms for complexes comprising hsp70, 50 to 5000 micrograms for hsp90, and 10 to 600 micrograms for gp96.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

ADOPTIVE IMMUNOTHERAPY UTILIZING HEAT SHOCK/STRESS PROTEIN-PEPTIDE COMPLEXES FOR PREVENTION/TREATMENT OF CANCER OR INFECTIOUS DISEASES

5 This invention was made with government support under grant numbers CA44786 and CA64394 awarded by the National Institutes of Health. The government has certain rights in the invention.

10 1. INTRODUCTION

 The present invention relates to methods and compositions for the prevention and treatment of infectious diseases, primary and metastatic neoplastic diseases, including, but not limited to human sarcomas and carcinomas.

15 In the practice of the invention, compositions of complexes of heat shock/stress proteins (hsps) including, but not limited to, hsp70, hsp90, gp96 alone or in combination with each other, noncovalently bound to antigenic molecules, are administered in combination with the administration of

20 antigen presenting cells sensitized with an hsp-antigenic molecule noncovalent complex to augment the immune response to genotoxic and nongenotoxic factors, tumors and infectious agents.

25 2. BACKGROUND OF THE INVENTION

 The era of tumor immunology began with experiments by Prehn and Main, who showed that antigens on the methylcholanthrene (MCA)-induced sarcomas were tumor specific in that transplantation assays could not detect these

30 antigens in normal tissue of the mice (Prehn, R.T., et al., 1957, *J. Natl. Cancer Inst.* 18:769-778). This notion was confirmed by further experiments demonstrating that tumor specific resistance against MCA-induced tumors can be elicited in the autochthonous host, that is, the mouse in

35 which the tumor originated (Klein, G., et al., 1960, *Cancer Res.* 20:1561-1572).

In subsequent studies, tumor specific antigens were also found on tumors induced with other chemical or physical carcinogens or on spontaneous tumors (Kripke, M.L., 1974, *J. Natl. Cancer Inst.* 53:1333-1336; Vaage, J., 1968, *Cancer Res.* 28:2477-2483; Carswell, E.A., et al., 1970, *J. Natl. Cancer Inst.* 44:1281-1288). Since these studies used protective immunity against the growth of transplanted tumors as the criterion for tumor specific antigens, these antigens are also commonly referred to as "tumor specific transplantation antigens" or "tumor specific rejection antigens." Several factors can greatly influence the immunogenicity of the tumor induced, including, for example, the specific type of carcinogen involved, immunocompetence of the host and latency period (Old, L.J., et al., 1962, *Ann. N.Y. Acad. Sci.* 101:80-106; Bartlett, G.L., 1972, *J. Natl. Cancer Inst.* 49:493-504).

Most, if not all, carcinogens are mutagens which may cause mutation, leading to the expression of tumor specific antigens (Ames, B.N., 1979, *Science* 204:587-593; Weisburger, J.H., et al., 1981, *Science* 214:401-407). Some carcinogens are immunosuppressive (Malmgren, R.A., et al., 1952, *Proc. Soc. Exp. Biol. Med.* 79:484-488). Experimental evidence suggests that there is a constant inverse correlation between immunogenicity of a tumor and latency period (time between exposure to carcinogen and tumor appearance) (Old, L.J., et al., 1962, *Ann. N.Y. Acad. Sci.* 101:80-106; and Bartlett, G.L., 1972, *J. Natl. Cancer Inst.* 49:493-504). Other studies have revealed the existence of tumor specific antigens that do not lead to rejection, but, nevertheless, can potentially stimulate specific immune responses (Roitt, I., Brostoff, J and Male, D., 1993, *Immunology*, 3rd ed., Mosby, St. Louis, pps. 17.1-17.12).

2.1. Tumor-Specific Immunogenicities of Heat Shock/Stress Proteins hsp70, hsp90 and gp96

Srivastava et al. demonstrated immune response to methylcholanthrene-induced sarcomas of inbred mice (1988, *Immunol. Today* 9:78-83). In these studies it was found that

the molecules responsible for the individually distinct immunogenicity of these tumors were identified as cell-surface glycoproteins of 96kDa (gp96) and intracellular proteins of 84 to 86kDa (Srivastava, P.K., et al., 1986, *Proc. Natl. Acad. Sci. USA* 83:3407-3411; Ullrich, S.J., et al., 1986, *Proc. Natl. Acad. Sci. USA* 83:3121-3125. Immunization of mice with gp96 or p84/86 isolated from a particular tumor rendered the mice immune to that particular tumor, but not to antigenically distinct tumors. Isolation and characterization of genes encoding gp96 and p84/86 revealed significant homology between them, and showed that gp96 and p84/86 were, respectively, the endoplasmic reticular and cytosolic counterparts of the same heat shock proteins (Srivastava, P.K., et al., 1988, *Immunogenetics* 28:205-207; Srivastava, P.K., et al., 1991, *Curr. Top. Microbiol. Immunol.* 167:109-123). Further, hsp70 was shown to elicit immunity to the tumor from which it was isolated but not to antigenically distinct tumors. However, hsp70 depleted of peptides was found to lose its immunogenic activity (Udono, M., and Srivastava, P.K., 1993, *J. Exp. Med.* 178:1391-1396). These observations suggested that the heat shock proteins are not immunogenic per se, but are carriers of antigenic peptides that elicit specific immunity to cancers (Srivastava, P.K., 1993, *Adv. Cancer Res.* 62:153-177).

25

2.2. Pathobiology of Cancer

Cancer is characterized primarily by an increase in the number of abnormal cells derived from a given normal tissue, invasion of adjacent tissues by these abnormal cells, and lymphatic or blood-borne spread of malignant cells to regional lymph nodes and to distant sites (metastasis). Clinical data and molecular biologic studies indicate that cancer is a multistep process that begins with minor preneoplastic changes, which may under certain conditions progress to neoplasia.

Pre-malignant abnormal cell growth is exemplified by hyperplasia, metaplasia, or most particularly, dysplasia (for

review of such abnormal growth conditions, see Robbins and Angell, 1976, *Basic Pathology*, 2d Ed., W.B. Saunders Co., Philadelphia, pp. 68-79.) Hyperplasia is a form of controlled cell proliferation involving an increase in cell
5 number in a tissue or organ, without significant alteration in structure or function. As but one example, endometrial hyperplasia often precedes endometrial cancer. Metaplasia is a form of controlled cell growth in which one type of adult or fully differentiated cell substitutes for another type of
10 adult cell. Metaplasia can occur in epithelial or connective tissue cells. Atypical metaplasia involves a somewhat disorderly metaplastic epithelium. Dysplasia is frequently a forerunner of cancer, and is found mainly in the epithelia; it is the most disorderly form of non-neoplastic cell growth,
15 involving a loss in individual cell uniformity and in the architectural orientation of cells. Dysplastic cells often have abnormally large, deeply stained nuclei, and exhibit pleomorphism. Dysplasia characteristically occurs where there exists chronic irritation or inflammation, and is often
20 found in the cervix, respiratory passages, oral cavity, and gall bladder.

The neoplastic lesion may evolve clonally and develop an increasing capacity for invasion, growth, metastasis, and heterogeneity, especially under conditions in which the
25 neoplastic cells escape the host's immune surveillance (Roitt, I., Brostoff, J and Kale, D., 1993, *Immunology*, 3rd ed., Mosby, St. Louis, pps. 17.1-17.12).

2.3. Immunotherapy

30 Four basic cell types whose function has been associated with antitumor cell immunity and the elimination of tumor cells from the body are: i) B-lymphocytes which secrete immunoglobulins into the blood plasma for identifying and labeling the nonself invader cells; ii) monocytes which
35 secrete the complement proteins which are responsible for lysing and processing the immunoglobulin-coated target invader cells; iii) natural killer lymphocytes having two

mechanisms for the destruction of tumor cells-antibody-dependent cellular cytotoxicity and natural killing; and iv) T-lymphocytes possessing antigen-specific receptors and each T-lymphocyte clone having the capacity to recognize a tumor cell carrying complementary marker molecules (Schreiber, H., 1989, in *Fundamental Immunology* (ed). W.E. Paul, pp. 923-955).

Several factors can influence the immunogenicity of tumors induced. These factors include dose of carcinogen, immunocompetence of the host, and latency period. Immunocompetence of the host during the period of cancer induction and development can allow the host to respond to immunogenic tumor cells. This may prevent the outgrowth of these cells or select far less immunogenic escape variants that have lost their respective rejection antigen.

Conversely, immunosuppression or immune deficiency of the host during carcinogenesis or tumorigenesis may allow growth of highly immunogenic tumors (Schreiber, H., 1989, in *Fundamental Immunology* (ed). W.E. Paul, pp. 923-955).

Three major types of cancer immunotherapy are currently being explored: i) adoptive cellular immunotherapy, ii) in vivo manipulation of patient plasma to remove blocking factors or add tumoricidal factors, and iii) in vivo administration of biological response modifiers (e.g., interferons (IFN; IFN-alpha and IFN-gamma), interleukins (IL; IL-2, IL-4 and IL-6), colony-stimulating factors, tumor necrosis factor (TNF), monoclonal antibodies and other immunopotentiating agents, such as corynebacterium parvum (C. parvum) (Kopp, W.C., et al., 1994, *Cancer Chemotherapy and Biol. Response Modifiers* 15:226-286). There is little doubt that immunotherapy of cancer as it stands is falling short of the hopes invested in it. Although numerous immunotherapeutic approaches have been tested, few of these procedures have proved to be effective as the sole or even as an adjunct form of cancer prevention and treatment.

2.3.1. Adoptive Cellular Immunotherapy

Adoptive immunotherapy of cancer refers to a therapeutic approach in which immune cells with an antitumor reactivity are administered to a tumor-bearing host, with the aim that the cells mediate either directly or indirectly, the regression of an established tumor. Transfusion of lymphocytes, particularly T lymphocytes, falls into this category and investigators at the National Cancer Institute (NCI) have used autologous reinfusion of peripheral blood lymphocytes or tumor-infiltrating lymphocytes (TIL), T cell cultures from biopsies of subcutaneous lymph nodes, to treat several human cancers (Rosenberg, S.A., U.S. Patent No. 4,690,914, issued September 1, 1987; Rosenberg, S.A., et al., 1988, *N. England J. Med.* 319:1676-1680). For example, TIL expanded *in vitro* in the presence of interleukin (IL)-2 have been adoptively transferred to cancer patients, resulting in tumor regression in select patients with metastatic melanoma. Melanoma TIL grown in IL-2 have been identified as activated T lymphocytes CD3⁺ HLA-DR⁺, which are predominantly CD8⁺ cells with unique *in vitro* antitumor properties. Many long-term melanoma TIL cultures lyse autologous tumors in a specific MHC class I- and T cell antigen receptor dependent manner (Topalian, S.L., et al., 1989, *J. Immunol.* 142:3714). However, studies of TIL derived from other types of tumors have revealed only scant evidence for cytolytic or proliferative antitumor immune specificity (Topalian, S.L. et al., 1990, in *Important Advances in Oncology*, V.T. DeVita, S.A. Hellman and S.A. Rosenberg, eds. J.B. Lippincott, Philadelphia, pp. 19-41). In addition, the toxicity of the high-dose IL-2 + activated lymphocyte treatment advocated by the NCI group has been considerable, including high fevers, severe rigors, hypotension, damage to the endothelial wall due to capillary leak syndrome, and various adverse cardiac events such as arrhythmias and myocardial infarction (Rosenberg S.A., et al., 1988, *N. England J. Med.* 319:1676-1680).

2.3.2. Interleukins (IL-2, IL-4 and IL-6)

IL-2 has significant antitumor activity in a small percentage of patients with renal cell carcinoma and melanoma. Investigators continue to search for IL-2 based regimens that will increase the response rates in IL-2 responsive tumors, but, for the most part, have neither defined new indications nor settled fundamental issues, such as whether dose intensity is important in IL-2 therapy (Kopp, W.C., et al., 1994, *Cancer Chemotherapy and Biol. Response* 10 *Modifiers* 15:226-286). Numerous reports have documented IL-2 associated toxicity involving increased nitrate levels and the syndrome of vascular leak and hypotension, analogous to septic shock. In addition, an increased incidence of nonopportunistic bacterial infections and autoimmune complications are frequently accompanied by the antitumor response of IL-2 (Kopp, W.C., et al., 1994, *Cancer Chemotherapy and Biol. Response Modifiers* 15:226-286).

IL-4 and IL-6 are also being tested as antitumor agents either directly or through immunomodulating mechanisms. Dose-limiting toxicities have been observed with both agents in Phase I clinical trials (Gilleece, M.H., et al., 1992, *Br. J. Cancer* 66:204-210, Weber, J., et al., 1993, *J. Clin. Oncol.* 11:499-506).

2.3.3. Tumor Necrosis Factor

The toxicity of systemically administered TNF seriously limits its use for the treatment of cancer. TNF has been most effective when used for regional therapy, in which measures, such as limb isolation for perfusion, are taken to limit the systemic dose and hence the toxicity of TNF. Dose-limiting toxicity of TNF consist of thrombocytopenia, headache, confusion and hypotension (Mittleman, A., et al., 1992, *Inv. New Drugs* 10:183-190).

2.3.4. Interferons

The activity of IFN- α has been described as being modest in a number of malignancies, including renal cell carcinoma,

melanoma, hairy cell leukemia low-grade non-Hodgkin's lymphoma, and others. Higher doses of IFN- α are usually associated with higher response rates in some malignancies, but also cause more toxicity. In addition, more and more reports indicate that relapses after successful interferon therapy coincide with formation of neutralizing antibodies against interferon (Ouesada, J.R., et al., 1987, *J. Interferon Res.* 67:678.

10 3. SUMMARY OF THE INVENTION

The methods of the invention comprise methods of eliciting an immune response in an individual in whom the treatment or prevention of cancer or infectious disease is desired by administering a composition comprising an effective amount of a complex in which the complex consists essentially of heat shock protein(s) (hsp(s)) noncovalently bound to antigenic molecule(s) in combination with adoptive immunotherapy involving administration of antigen presenting cells that have been sensitized *in vitro* with complexes of hsp(s) noncovalently bound to antigenic molecule(s). Such adoptive immunotherapy in combination with administration of hsp-antigenic molecule complexes is believed to exhibit a synergistic effect in eliciting an immune response.

The amounts of the complex that are administered are within ranges of effective dosages, discovered by the present inventor to be effective. The lower limit of the ranges is surprisingly smaller than those amounts predicted to be effective by extrapolation by prior art methods from dosages used in animal studies. In a preferred embodiment, the complex is autologous to the individual; that is, the complex is isolated from the cancer cells of the individual himself (e.g., preferably prepared from tumor biopsies of the patient). Alternatively, the hsp and or the antigenic molecule can be isolated from the individual or from others or by recombinant production methods using a cloned hsp originally derived from the individual or from others.

"Antigenic molecule" as used herein refers to the peptides

with which the hsps are endogenously associated *in vivo* (e.g., in precancerous or cancerous tissue), as well as exogenous antigens/immunogens (i.e., with which the hsps are not complexed *in vivo*) or antigenic/immunogenic fragments and derivatives thereof. Such exogenous antigens and fragments and derivatives (both peptide and non-peptide) thereof for use in complexing with hsps, can be selected from among those known in the art, as well as those readily identified by standard immunoassays known in the art by detecting the ability to bind antibody or MHC molecules (antigenicity) or generate immune response (immunogenicity).

The methods for adoptive immunotherapy of cancer and infectious diseases have the goal of enhancing the host's immunocompetence and activity of immune effector cells. Adoptive immunotherapy with macrophages and/or other antigen-presenting cells (APC), for example, dendritic cells or B cells (B lymphocytes), that have been sensitized *in vitro* with noncovalent complexes of hsp noncovalently bound to an antigenic molecule, induces specific immunity to tumor cells and/or antigenic components, promoting regression of the tumor mass or treatment of immunological disorders or infectious diseases, as the case may be.

In a specific embodiment, the present invention relates to methods and compositions for prevention and treatment of primary and metastatic neoplastic diseases.

Specific therapeutic regimens, pharmaceutical compositions, and kits are provided by the invention. The preferred dosages of the hsp-antigenic molecule complexes are less than those based on body weight or surface area of the patient. The present inventor has discovered that a dosage substantially equivalent to or smaller than that seen to be effective in smaller non-human mammals (e.g., mice) is effective for human administration, optionally subject to a correction factor not exceeding a fifty fold increase, based on the relative lymph node sizes in such mammals and in humans.

The mode of administration of the hsp-antigenic molecule complexes includes but is not limited to, e.g., subcutaneously, intramuscularly, intravenously, intraperitoneally, intradermally or mucosally. The present
5 inventor has discovered that effective intradermal dosages of the hsp-antigenic molecule complexes are about tenfold smaller even than the surprisingly small doses effective in subcutaneous administration in humans. (See U.S. patent application Serial No. 08/527,391, filed September 13, 1995,
10 which is incorporated by reference herein in its entirety, and U.S. patent application entitled "Prevention and Treatment of Primary and Metastatic Neoplastic Diseases and Infectious Diseases With Heat Shock/Stress Protein-Peptide Complexes" by P. Srivastava and R. Chandawarkar, filed on
15 even date herewith, which is incorporated by reference herein in its entirety.) Pharmaceutical formulations are provided comprising compositions of complexes of antigenic molecules and heat shock/stress proteins, including but not limited to hsp70, hsp90, gp96 either alone or in combination, to be
20 administered before, concurrently with, or after the administration of the hsp complex-sensitized APC. Specifically, interspecies dose-response equivalence for hsp noncovalently bound to antigenic molecules for a human intradermal or mucosal dose is estimated as the product of
25 the therapeutic dosage observed in mice and a single scaling ratio, not exceeding a fifty fold increase.

The present invention encompasses methods for prevention and treatment of cancer by enhancing the host's immune competence and activity of immune effector cells.
30 Furthermore, the invention provides methods for evaluating the efficacy of drugs in enhancing immune responses for treatment and monitoring the progress of patients participating in clinical trials for the treatment of primary and metastatic neoplastic diseases.
35 Immunotherapy using the therapeutic regimens of the invention, by administering such complexes of heat shock/stress proteins noncovalently bound to antigenic

molecules in combination with adoptive immunotherapy, can induce specific immunity to tumor cells, and can lead to regression of the tumor mass. Cancers which are responsive to specific immunotherapy by the methods of the invention
5 include but are not limited to human sarcomas and carcinomas. In a specific embodiment, the hsp-antigenic molecule complexes are allogeneic to the patient; in a preferred embodiment, the hsp-antigenic molecule complexes are autologous to (derived from) the patient to whom they are
10 administered.

Particular compositions of the invention and their properties are described in the sections and subsections which follow. A preferred composition for both administration to the patient and for use in sensitizing APC
15 to be administered to the patient comprises hsp-peptide complexes isolated from the tumor biopsy of the patient to whom the composition is to be administered. Such a composition that comprises hsp70, hsp90 and/or gp96 demonstrates strong inhibition of a variety of tumors in
20 mammals. Preferred compositions comprising hsp70, hsp90 and/or gp96 which preferably exhibit no toxicity when administered to human subjects are also described.

In another embodiment, the methods further optionally comprise administering biological response modifiers, e.g.,
25 IFN- α , IFN- γ , IL-2, IL-4, IL-6, TNF, or other cytokine growth factors affecting the immune cells, in combination with the hsp complexes.

In addition to cancer therapy, the administration of complexes of hsps noncovalently bound to antigenic molecules
30 in combination with adoptive immunotherapy methods of the invention can be utilized for the prevention of a variety of cancers, e.g., in individuals who are predisposed as a result of familial history or in individuals with an enhanced risk to cancer due to environmental factors.

35 The Examples presented in Sections 5 and 6 below, detail the use according to the methods of the invention of hsp-peptide complexes in combination with adoptive immunotherapy

in cancer immunotherapy in human patients suffering from advanced colon and liver cancer.

4. DETAILED DESCRIPTION OF THE INVENTION

5 Methods and compositions for the prevention and treatment of primary and metastatic neoplastic diseases and infectious diseases and for eliciting an immune response in a human individual, are described. The invention is based on the effect of stimulating the immune system by administration
10 of compositions comprising complexes of hsps noncovalently bound to antigenic molecules in combination with adoptive immunotherapy comprising the administration of antigen presenting cells (APC) sensitized *in vitro* with complexes of hsps noncovalently bound to antigenic molecules. It is
15 believed that the effect of such combination therapy may be synergistic.

"Antigenic molecule" as used herein refers to the peptides with which the hsps are endogenously associated *in vivo* (e.g., in infected cells or precancerous or cancerous
20 tissue) as well as exogenous antigens/immunogens (i.e., with which the hsps are not complexed *in vivo*) or antigenic/immunogenic fragments and derivatives thereof.

The methods of the invention comprise methods of eliciting an immune response in an individual in whom the
25 treatment or prevention of infectious diseases or cancer is desired by administering a composition comprising an effective amount of a complex, in which the complex consists essentially of an hsp noncovalently bound to an antigenic molecule using any convenient mode of administration in
30 combination with the adoptive immunotherapy methods of the invention. Modes of administration include but are not limited to subcutaneously, intramuscularly, intravenously, intraperitoneally, intradermally or mucosally.

The APC can be selected from among those antigen
35 presenting cells known in the art, including but not limited to macrophages, dendritic cells, B lymphocytes, and a combination thereof, and are preferably macrophages. The hsp

complex-sensitized APC may be administered concurrently or before or after administration of the hsp-antigenic molecule complexes. Adoptive immunotherapy according to the invention allows activation of antigen presenting cells (APC) by
5 incubation *in vitro* with hsp-antigenic molecule complexes. Preferably, prior to use of the cells *in vivo*, measurement of reactivity against the tumor or infectious agent *in vitro* is done. This *in vitro* boost followed by clonal selection and/or expansion, and patient administration constitutes a
10 useful therapeutic/prophylactic strategy.

In a preferred embodiment, the hsp-antigenic molecule complex is autologous to the individual; that is, the complex is isolated from either the infected cells or the cancer cells or precancerous cells of the individual himself (e.g.,
15 preferably prepared from infected tissues or tumor biopsies of the patient). Alternatively, the complex is produced *in vitro* (e.g., wherein a complex with an exogenous antigenic molecule is desired). Alternatively, the hsp and/or the antigenic molecule can be isolated from the individual or
20 from others or made by recombinant production methods using a cloned hsp originally derived from the individual or from others. Exogenous antigens and fragments and derivatives (both peptide and non-peptide) thereof for use in complexing with hsps, can be selected from among those known in the art,
25 as well as those readily identified by standard immunoassays known in the art by the ability to bind antibody or MHC molecules (antigenicity) or generate immune response (immunogenicity). Complexes of hsps and antigenic molecules can be isolated from cancer or precancerous tissue of a
30 patient, or from a cancer cell line, or can be produced *in vitro* (as is necessary in the embodiment in which an exogenous antigen is used as the antigenic molecule). The hsp-antigenic molecule complex that is administered to the patient can be the same or different from the hsp-antigenic
35 molecule complex used to sensitize the APC that are administered to the patient. In a specific embodiment wherein the APC and complexes are administered concurrently,

the APC and purified hsp-antigenic molecule complexes can be present in a single composition, or different compositions, for administration.

The hsps of the present invention that can be used
5 include but are not limited to, hsp70, hsp90, gp96 alone or in combination. Preferably, the hsps are human hsps.

Heat shock proteins, which are also referred to interchangeably herein as stress proteins, useful in the practice of the instant invention can be selected from among
10 any cellular protein that satisfies any one of the following criteria. It is a protein whose intracellular concentration increases when a cell is exposed to a stressful stimuli, it is capable of binding other proteins or peptides, it is capable of releasing the bound proteins or peptides in the
15 presence of adenosine triphosphate (ATP) or low pH, or it is a protein showing at least 35% homology with any cellular protein having any of the above properties.

The first stress proteins to be identified were the heat shock proteins (hsps). As their name implies, hsps are
20 synthesized by a cell in response to heat shock. To date, three major families of hsp have been identified based on molecular weight. The families have been called hsp60, hsp70 and hsp90 where the numbers reflect the approximate molecular weight of the stress proteins in kilodaltons. Many members
25 of these families were found subsequently to be induced in response to other stressful stimuli including, but not limited to, nutrient deprivation, metabolic disruption, oxygen radicals, and infection with intracellular pathogens.

(See Welch, May 1993, *Scientific American* 56-64; Young, 1990, *Annu. Rev. Immunol.* 8:401-420; Craig, 1993, *Science* 260:1902-
30 1903; Gething, et al., 1992, *Nature* 355:33-45; and Lindquist, et al., 1988, *Annu. Rev. Genetics* 22:631-677), the disclosures of which are incorporated herein by reference.

It is contemplated that hsps/stress proteins belonging to all
35 of these three families can be used in the practice of the instant invention.

The major hsps can accumulate to very high levels in stressed cells, but they occur at low to moderate levels in cells that have not been stressed. For example, the highly inducible mammalian hsp70 is hardly detectable at normal
5 temperatures but becomes one of the most actively synthesized proteins in the cell upon heat shock (Welch, et al., 1985, *J. Cell. Biol.* 101:1198-1211). In contrast, hsp90 and hsp60 proteins are abundant at normal temperatures in most, but not all, mammalian cells and are further induced by heat (Lai, et
10 al., 1984, *Mol. Cell. Biol.* 4:2802-10; van Bergen en Henegouwen, et al., 1987, *Genes Dev.* 1:525-31).

Heat shock proteins are among the most highly conserved proteins in existence. For example, DnaK, the hsp70 from *E. coli* has about 50% amino acid sequence identity with hsp70
15 proteins from excoriates (Bardwell, et al., 1984, *Proc. Natl. Acad. Sci.* 81:848-852). The hsp60 and hsp90 families also show similarly high levels of intrafamilies conservation (Hickey, et al., 1989, *Mol. Cell. Biol.* 9:2615-2626; Jindal, 1989, *Mol. Cell. Biol.* 9:2279-2283). In addition, it has
20 been discovered that the hsp60, hsp70 and hsp90 families are composed of proteins that are related to the stress proteins in sequence, for example, having greater than 35% amino acid identity, but whose expression levels are not altered by stress. Therefore it is contemplated that the definition of
25 heat shock protein or stress protein, as used herein, embraces other proteins, muteins, analogs, and variants thereof having at least 35% to 55%, preferably 55% to 75%, and most preferably 75% to 85% amino acid identity with members of the three families whose expression levels in a
30 cell are enhanced in response to a stressful stimulus. The purification of stress proteins belonging to these three families is described below.

The immunogenic hsp-peptide complexes of the invention may include any complex containing an hsp and a peptide that
35 is capable of inducing an immune response in a mammal. The peptides are preferably noncovalently associated with the hsp. Preferred complexes may include, but are not limited

to, hsp60-peptide, hsp70-peptide and hsp90-peptide complexes. For example, an hsp called gp96 which is present in the endoplasmic reticulum of eukaryotic cells and is related to the cytoplasmic hsp90's can be used to generate an effective
5 vaccine containing a gp96-peptide complex.

Although the hsps can be allogeneic to the patient, in a preferred embodiment, the hsps are autologous to (derived from) the patient to whom they are administered. The hsps and/or antigenic molecules can be purified from natural
10 sources, chemically synthesized, or recombinantly produced.

The invention provides combinations of compositions which enhance the immunocompetence of the host individual and elicit specific immunity against infectious agents or specific immunity against preneoplastic and neoplastic cells.
15 The therapeutic regimens and pharmaceutical compositions of the invention are described below. These compositions have the capacity to prevent the onset and progression of infectious diseases and prevent the development of tumor cells and to inhibit the growth and progression of tumor
20 cells indicating that such compositions can induce specific immunity in infectious diseases and cancer immunotherapy.

Hsps appear to induce an inflammatory reaction at the tumor site and ultimately cause a regression of the tumor burden in the cancer patients treated. Cancers which can be
25 treated with complexes of hsps noncovalently bound to antigenic molecules include, but are not limited to, human sarcomas and carcinomas. Human sarcomas and carcinomas are also responsive to adoptive immunotherapy by the hsp complex-sensitized macrophages and/or APC.

30 Accordingly, the invention provides methods of preventing and treating cancer in an individual comprising administering hsp-antigenic molecule complexes, and APC sensitized by such complexes, which stimulates the immunocompetence of the host individual and elicits specific
35 immunity against the preneoplastic and/or neoplastic cells.

As used herein, "preneoplastic" cell refers to a cell which is in transition from a normal to a neoplastic form; and

morphological evidence, increasingly supported by molecular biologic studies, indicates that preneoplasia progresses through multiple steps. Non-neoplastic cell growth commonly consists of hyperplasia, metaplasia, or most particularly, 5 dysplasia (for review of such abnormal growth conditions (See Robbins and Angell, 1976, *Basic Pathology*, 2d Ed., W.B. Saunders Co., Philadelphia, pp. 68-79). Hyperplasia is a form of controlled cell proliferation involving an increase in cell number in a tissue or organ, without significant 10 alteration in structure or function. As but one example, endometrial hyperplasia often precedes endometrial cancer. Metaplasia is a form of controlled cell growth in which one type of adult or fully differentiated cell substitutes for another type of adult cell. Metaplasia can occur in 15 epithelial or connective tissue cells. Atypical metaplasia involves a somewhat disorderly metaplastic epithelium. Dysplasia is frequently a forerunner of cancer, and is found mainly in the epithelia; it is the most disorderly form of non-neoplastic cell growth, involving a loss in individual 20 cell uniformity and in the architectural orientation of cells. Dysplastic cells often have abnormally large, deeply stained nuclei, and exhibit pleomorphism. Dysplasia characteristically occurs where there exists chronic irritation or inflammation, and is often found in the cervix, 25 respiratory passages, oral cavity, and gall bladder. Although preneoplastic lesions may progress to neoplasia, they may also remain stable for long periods and may even regress, particularly if the inciting agent is removed or if the lesion succumbs to an immunological attack by its host.

30 The present inventor has discovered that a dosage substantially equivalent to or smaller than that seen to be effective in smaller non-human animals (e.g., mice) is effective for human administration, such as described in Section 4.1, below.

35 The therapeutic regimens and pharmaceutical compositions of the invention may be used with additional immune response enhancers or biological response modifiers including, but not

limited to, the cytokines IFN- α , IFN- γ , IL-2, IL-4, IL-6, TNF, or other cytokine affecting immune cells. In accordance with this aspect of the invention, the complexes of the hsp and antigenic molecule are administered in combination
5 therapy with one or more of these cytokines.

The invention further relates to administration of complexes of hsp-antigenic molecules in combination with adoptive immunotherapy to individuals at enhanced risk of cancer due to familial history or environmental risk factors.

10

**4.1. Compositions Comprising Hsp-Peptide
Complexes, for Eliciting Immune Responses
to Cancer or Infectious Diseases, and for
In Vitro Sensitization of APC**

The compositions comprising hsp noncovalently bound to
15 antigenic molecules are administered to elicit an effective specific immune response to the complexed antigenic molecules (and not to the hsp). In accordance with the methods described herein, the hsp-antigenic molecule complexes are preferably purified in the range of 60 to 100 percent of the
20 total mg protein, or at least 70%, 80% or 90% of the total mg protein. In another embodiment, the hsp-antigenic molecule complexes are purified to apparent homogeneity, as assayed by sodium dodecyl sulfate-polyacrylamide gel electrophoresis.

In a preferred embodiment, non-covalent complexes of
25 hsp70, hsp90 and gp96 with peptides are prepared and purified postoperatively from tumor cells obtained from the cancer patient.

In accordance with the methods described herein, immunogenic or antigenic peptides that are endogenously
30 complexed to hsps or MHC antigens can be used as antigenic molecules. For example, such peptides may be prepared that stimulate cytotoxic T cell responses against different tumor antigens (e.g., tyrosinase, gp100, melan-A, gp75, mucins, etc.) and viral proteins including, but not limited to,
35 proteins of immunodeficiency virus type I (HIV-I), human immunodeficiency virus type II (HIV-II), hepatitis type A, hepatitis type B, hepatitis type C, influenza, Varicella,

adenovirus, herpes simplex type I (HSV-I), herpes simplex type II (HSV-II), rinderpest, rhinovirus, echovirus, rotavirus, respiratory syncytial virus, papilloma virus, papova virus, cytomegalovirus, echinovirus, arbovirus, 5 huntavirus, coxsackie virus, mumps virus, measles virus, rubella virus and polio virus. In the embodiment wherein the antigenic molecules are peptides noncovalently complexed to hsps *in vivo*, the complexes can be isolated from cells, or alternatively, produced *in vitro* from purified preparations 10 each of hsps and antigenic molecules.

In another specific embodiment, antigens of cancers (e.g., tumors) or infectious agents (e.g., viral antigen, bacterial antigens, etc.) can be obtained by purification from natural sources, by chemical synthesis, or 15 recombinantly, and, through *in vitro* procedures such as that described below, noncovalently complexed to hsps.

In an embodiment wherein the hsp-antigenic molecule complex to be used is a complex that is produced *in vivo* in cells, exemplary purification procedures such as described in 20 Sections 4.1.1-4.1.3 below can be employed. Alternatively, in an embodiment wherein one wishes to use antigenic molecules by complexing to hsps *in vitro*, hsps can be purified for such use from the endogenous hsp-peptide complexes in the presence of ATP or low pH (or chemically 25 synthesized or recombinantly produced). The protocols described herein may be used to isolate hsp-peptide complexes, or the hsps alone, from any eukaryotic cells for example, tissues, isolated cells, or immortalized eukaryote cell lines infected with a preselected intracellular 30 pathogen, tumor cells or tumor cell lines.

4.1.1. Preparation and Purification of Hsp70-peptide Complexes

The purification of hsp70-peptide complexes has been described previously, see, for example, Udono et al., 1993, *J. Exp. Med.* 178:1391-1396. A procedure that may be used, presented by way of example but not limitation, is as follows:

Initially, tumor cells are suspended in 3 volumes of 1X Lysis buffer consisting of 5mM sodium phosphate buffer, pH 7, 150mM NaCl, 2mM CaCl₂, 2mM MgCl₂, and 1mM phenyl methyl sulfonyl fluoride (PMSF). Then, the pellet is sonicated, on ice, until >99% cells are lysed as determined by microscopic examination. As an alternative to sonication, the cells may be lysed by mechanical shearing and in this approach the cells typically are resuspended in 30mM sodium bicarbonate pH 7.5, 1mM PMSF, incubated on ice for 20 minutes and then homogenized in a Dounce homogenizer until >95% cells are lysed.

Then the lysate is centrifuged at 1,000g for 10 minutes to remove unbroken cells, nuclei and other cellular debris. The resulting supernatant is recentrifuged at 100,000g for 90 minutes, the supernatant harvested and then mixed with Con A Sepharose equilibrated with phosphate buffered saline (PBS) containing 2mM Ca²⁺ and 2mM Mg²⁺. When the cells are lysed by mechanical shearing the supernatant is diluted with an equal volume of 2X lysis buffer prior to mixing with Con A Sepharose. The supernatant is then allowed to bind to the Con A Sepharose for 2-3 hours at 4°C. The material that fails to bind is harvested and dialyzed for 36 hours (three times, 100 volumes each time) against 10mM Tris-Acetate pH 7.5, 0.1mM EDTA, 10mM NaCl, 1mM PMSF. Then the dialyzate is centrifuged at 17,000 rpm (Sorvall SS34 rotor) for 20 minutes. Then the resulting supernatant is harvested and applied to a Mono Q FPLC column equilibrated in 20mM Tris-Acetate pH 7.5, 20mM NaCl, 0.1mM EDTA and 15mM 2-mercaptoethanol. The column is then developed with a 20mM to 500mM NaCl gradient and then eluted fractions fractionated

by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) and characterized by immunoblotting using an appropriate anti-hsp70 antibody (such as from clone N27F3-4, from StressGen).

5 Fractions strongly immunoreactive with the anti-hsp70 antibody are pooled and the hsp70-peptide complexes precipitated with ammonium sulfate; specifically with a 50%-70% ammonium sulfate cut. The resulting precipitate is then harvested by centrifugation at 17,000 rpm (SS34 Sorvall
10 rotor) and washed with 70% ammonium sulfate. The washed precipitate is then solubilized and any residual ammonium sulfate removed by gel filtration on a Sephadex^R G25 column (Pharmacia). If necessary the hsp70 preparation thus obtained can be repurified through the Mono Q FPLC Column as
15 described above.

The hsp70-peptide complex can be purified to apparent homogeneity using this method. Typically 1 mg of hsp70-peptide complex can be purified from 1 g of cells/tissue.

An improved method for purification of hsp70-peptide
20 complexes comprises contacting cellular proteins with ADP or a nonhydrolyzable analog of ATP affixed to a solid substrate, such that hsp70 in the lysate can bind to the ADP or nonhydrolyzable ATP analog, and eluting the bound hsp70. A preferred method uses column chromatography with ADP affixed
25 to a solid substratum (e.g., ADP-agarose). The resulting hsp70 preparations are higher in purity and devoid of contaminating peptides. The hsp70 yields are also increased significantly by about more than 10 fold. Alternatively, chromatography with nonhydrolyzable analogs of ATP, instead
30 of ADP, can be used for purification of hsp70-peptide complexes. By way of example but not limitation, purification of hsp70-peptide complexes by ADP-agarose chromatography can be carried out as follows:

Meth A sarcoma cells (500 million cells) are homogenized
35 in hypotonic buffer and the lysate is centrifuged at 100,000 g for 90 minutes at 4°C. The supernatant is applied to an ADP-agarose column. The column is washed in buffer and is

eluted with 5 column volumes of 3 mM ADP. The hsp70-peptide complexes elute in fractions 2 through 10 of the total 15 fractions which elute. The eluted fractions are analyzed by SDS-PAGE. The hsp70-peptide complexes can be purified to 5 apparent homogeneity using this procedure.

4.1.2. Preparation and Purification of Hsp90-peptide Complexes

A procedure that can be used, presented by way of example and not limitation, is as follows:

Initially, tumor cells are suspended in 3 volumes of 1X Lysis buffer consisting of 5mM sodium phosphate buffer (pH7), 150mM NaCl, 2mM CaCl₂, 2mM MgCl₂ and 1mM phenyl methyl sulfonyl fluoride (PMSF). Then, the pellet is sonicated, on ice, until >99% cells are lysed as determined by microscopic examination. As an alternative to sonication, the cells may be lysed by mechanical shearing and in this approach the cells typically are resuspended in 30mM sodium bicarbonate pH 7.5, 1mM PMSF, incubated on ice for 20 minutes and then homogenized in a Dounce homogenizer until >95% cells are lysed.

Then the lysate is centrifuged at 1,000g for 10 minutes to remove unbroken cells, nuclei and other cellular debris. The resulting supernatant is recentrifuged at 100,000g for 90 minutes, the supernatant harvested and then mixed with Con A Sepharose equilibrated with PBS containing 2mM Ca²⁺ and 2mM Mg²⁺. When the cells are lysed by mechanical shearing the supernatant is diluted with an equal volume of 2X Lysis buffer prior to mixing with Con A Sepharose. The supernatant is then allowed to bind to the Con A Sepharose for 2-3 hours at 4°C. The material that fails to bind is harvested and dialyzed for 36 hours (three times, 100 volumes each time) against 10mM Tris-Acetate pH 7.5, 0.1mM EDTA, 10mM NaCl, 1mM PMSF. Then the dialyzate is centrifuged at 17,000 rpm (Sorvall SS34 rotor) for 20 minutes. Then the resulting supernatant is harvested and applied to a Mono Q FPLC column

equilibrated with lysis buffer. The proteins are then eluted with a salt gradient of 200mM to 600mM NaCl.

The eluted fractions are fractionated by SDS-PAGE and fractions containing the hsp90-peptide complexes identified by immunoblotting using an anti-hsp90 antibody such as 3G3 (Affinity Bioreagents). Hsp90-peptide complexes can be purified to apparent homogeneity using this procedure. Typically, 150-200 μ g of hsp90-peptide complex can be purified from 1g of cells/tissue.

10

4.1.3. Preparation and Purification of gp96-peptide Complexes

A procedure that can be used, presented by way of example and not limitation, is as follows:

15 A pellet of tumors is resuspended in 3 volumes of buffer consisting of 30mM sodium bicarbonate buffer (pH 7.5) and 1mM PMSF and the cells allowed to swell on ice 20 minutes. The cell pellet is then homogenized in a Dounce homogenizer (the appropriate clearance of the homogenizer will vary according to each cell type) on ice until >95% cells are lysed.

20 The lysate is centrifuged at 1,000g for 10 minutes to remove unbroken cells, nuclei and other debris. The supernatant from this centrifugation step is then recentrifuged at 100,000g for 90 minutes. The gp96-peptide complex can be purified either from the 100,000 pellet or 25 from the supernatant.

When purified from the supernatant, the supernatant is diluted with equal volume of 2X lysis buffer and the supernatant mixed for 2-3 hours at 4°C with Con A Sepharose equilibrated with PBS containing 2mM Ca^{2+} and 2mM Mg^{2+} . Then, 30 the slurry is packed into a column and washed with 1X lysis buffer until the OD_{280} drops to baseline. Then, the column is washed with 1/3 column bed volume of 10% α -methyl mannoside (α -MM) dissolved in PBS containing 2mM Ca^{2+} and 2mM Mg^{2+} , the column sealed with a piece of parafilm, and incubated at 37°C 35 for 15 minutes. Then the column is cooled to room temperature and the parafilm removed from the bottom of the

column. Five column volumes of the α -MM buffer are applied to the column and the eluate analyzed by SDS-PAGE. Typically the resulting material is about 60-95% pure, however this depends upon the cell type and the tissue-to-lysis buffer ratio used. Then the sample is applied to a Mono Q FPLC column (Pharmacia) equilibrated with a buffer containing 5mM sodium phosphate, pH 7. The proteins are then eluted from the column with a 0-1M NaCl gradient and the gp96 fraction elutes between 400mM and 550mM NaCl.

- 10 The procedure, however, may be modified by two additional steps, used either alone or in combination, to consistently produce apparently homogeneous gp96-peptide complexes. One optional step involves an ammonium sulfate precipitation prior to the Con A purification step and the
15 other optional step involves DEAE-Sepharose purification after the Con A purification step but before the Mono Q FPLC step.

In the first optional step, described by way of example as follows, the supernatant resulting from the 100,000g
20 centrifugation step is brought to a final concentration of 50% ammonium sulfate by the addition of ammonium sulfate. The ammonium sulfate is added slowly while gently stirring the solution in a beaker placed in a tray of ice water. The solution is stirred from about 1/2 to 12 hours at 4°C and the
25 resulting solution centrifuged at 6,000 rpm (Sorvall SS34 rotor). The supernatant resulting from this step is removed, brought to 70% ammonium sulfate saturation by the addition of ammonium sulfate solution, and centrifuged at 6,000 rpm (Sorvall SS34 rotor). The resulting pellet from this step is
30 harvested and suspended in PBS containing 70% ammonium sulfate in order to rinse the pellet. This mixture is centrifuged at 6,000 rpm (Sorvall SS34 rotor) and the pellet dissolved in PBS containing 2mM Ca^{2+} and Mg^{2+} . Undissolved material is removed by a brief centrifugation at 15,000 rpm
35 (Sorvall SS34 rotor). Then, the solution is mixed with Con A Sepharose and the procedure followed as before.

In the second optional step, described by way of example as follows, the gp96 containing fractions eluted from the Con A column are pooled and the buffer exchanged for 5mM sodium phosphate buffer, pH 7, 300mM NaCl by dialysis, or preferably
5 by buffer exchange on a Sephadex G25 column. After buffer exchange, the solution is mixed with DEAE-Sepharose previously equilibrated with 5mM sodium phosphate buffer, pH 7, 300mM NaCl. The protein solution and the beads are mixed gently for 1 hour and poured into a column. Then, the column
10 is washed with 5mM sodium phosphate buffer, pH 7, 300mM NaCl, until the absorbance at 280nm drops to baseline. Then, the bound protein is eluted from the column with five volumes of 5mM sodium phosphate buffer, pH 7, 700mM NaCl. Protein containing fractions are pooled and diluted with 5mM sodium
15 phosphate buffer, pH 7 in order to lower the salt concentration to 175mM. The resulting material then is applied to the Mono Q FPLC column (Pharmacia) equilibrated with 5mM sodium phosphate buffer, pH 7 and the protein that binds to the Mono Q FPLC column (Pharmacia) is eluted as
20 described before.

It is appreciated, however, that one skilled in the art may assess, by routine experimentation, the benefit of incorporating the second optional step into the purification protocol. In addition, it is appreciated also that the
25 benefit of adding each of the optional steps will depend upon the source of the starting material.

When the gp96 fraction is isolated from the 100,000g pellet, the pellet is suspended in 5 volumes of PBS containing either 1% sodium deoxycholate or 1% octyl
30 glucopyranoside (but without the Mg^{2+} and Ca^{2+}) and incubated on ice for 1 hour. The suspension is centrifuged at 20,000g for 30 minutes and the resulting supernatant dialyzed against several changes of PBS (also without the Mg^{2+} and Ca^{2+}) to remove the detergent. The dialysate is centrifuged at
35 100,000g for 90 minutes, the supernatant harvested, and calcium and magnesium are added to the supernatant to give final concentrations of 2mM, respectively. Then the sample

is purified by either the unmodified or the modified method for isolating gp96-peptide complex from the 100,000g supernatant, see above.

The gp96-peptide complexes can be purified to apparent 5 homogeneity using this procedure. About 10-20µg of gp96 can be isolated from 1g cells/tissue.

Infectious Disease

In an alternative embodiment wherein it is desired to 10 treat a patient having an infectious disease, the above-described methods in Sections 4.1.1 - 4.1.3 are used to isolate hsp-peptide complexes from cells infected with an infectious organism, e.g., of a cell line or from a patient. Such infectious organisms include but are not limited to, 15 viruses, bacteria, protozoa, fungi, and parasites as described in detail in Section 4.1.4 below.

4.1.4. Isolation of Antigenic/Immunogenic Components

20 It has been found that antigenic peptides and/or components can be eluted from hsp-complexes either in the presence of ATP or low pH. These experimental conditions may be used to isolate peptides and/or antigenic components from cells which may contain potentially useful antigenic 25 determinants. Once isolated, the amino acid sequence of each antigenic peptide may be determined using conventional amino acid sequencing methodologies. Such antigenic molecules can then be produced by chemical synthesis or recombinant methods, purified, and complexed to hsps in vitro.

30 Similarly, it has been found that potentially immunogenic peptides may be eluted from MHC-peptide complexes using techniques well known in the art (Falk, K. et al., 1990 Nature 348:248-251; Elliott, T., et al., 1990, Nature 348:195-197; Falk, K., et al., 1991, Nature 351:290-296).

35 Thus, potentially immunogenic or antigenic peptides may be isolated from either endogenous stress protein-peptide complexes or endogenous MHC-peptide complexes for use

subsequently as antigenic molecules, by complexing *in vitro* to hsps. Exemplary protocols for isolating peptides and/or antigenic components from either of the these complexes are set forth below in Sections 4.1.4.1 and 4.1.4.2.

5

4.1.4.1. Peptides From Stress Protein-Peptide Complexes

Two methods may be used to elute the peptide from a stress protein-peptide complex. One approach involves incubating the stress protein-peptide complex in the presence of ATP. The other approach involves incubating the complexes in a low pH buffer.

Briefly the complex of interest is centrifuged through a Centricon 10 assembly (Millipore) to remove any low molecular weight material loosely associated with the complex. The large molecular weight fraction may be removed and analyzed by SDS-PAGE while the low molecular weight may be analyzed by HPLC as described below. In the ATP incubation protocol, the stress protein-peptide complex in the large molecular weight fraction is incubated with 10mM ATP for 30 minutes at room temperature. In the low pH protocol, acetic acid or trifluoroacetic acid (TFA) is added to the stress protein-peptide complex to give a final concentration of 10% (vol/vol) and the mixture incubated at room temperature or in a boiling water bath or any temperature in between, for 10 minutes (See, Van Bleek, et al., 1990, *Nature* 348:213-216; and Li, et al., 1993, *EMBO Journal* 12:3143-3151).

The resulting samples are centrifuged through a Centricon 10 assembly as mentioned previously. The high and low molecular weight fractions are recovered. The remaining large molecular weight stress protein-peptide complexes can be reincubated with ATP or low pH to remove any remaining peptides.

The resulting lower molecular weight fractions are pooled, concentrated by evaporation and dissolved in 0.1% TFA. The dissolved material is then fractionated by reverse phase high pressure liquid chromatography (HPLC) using for

example a VYDAC C18 reverse phase column equilibrated with 0.1% TFA. The bound material is then eluted at a flow rate of about 0.8 ml/min by developing the column with a linear gradient of 0 to 80% acetonitrile in 0.1% TFA. The elution
5 of the peptides can be monitored by OD₂₁₀ and the fractions containing the peptides collected.

4.1.4.2. Peptides from MHC-peptide Complexes

10 The isolation of potentially immunogenic peptides from MHC molecules is well known in the art and so is not described in detail herein (See, Falk, et al., 1990, *Nature* 348:248-251; Rotzsche, et al., 1990, *Nature* 348:252-254; Elliott, et al., 1990, *Nature* 348:191-197; Falk, et al.,
15 1991, *Nature* 351:290-296; Demotz, et al., 1989, *Nature* 343:682-684; Rotzsche, et al., 1990, *Science* 249:283-287), the disclosures of which are incorporated herein by reference.

Briefly, MHC-peptide complexes may be isolated by a
20 conventional immunoaffinity procedure. The peptides then may be eluted from the MHC-peptide complex by incubating the complexes in the presence of about 0.1% TFA in acetonitrile. The eluted peptides may be fractionated and purified by reverse phase HPLC, as before.

25 The amino acid sequences of the eluted peptides may be determined either by manual or automated amino acid sequencing techniques well known in the art. Once the amino acid sequence of a potentially protective peptide has been determined the peptide may be synthesized in any desired
30 amount using conventional peptide synthesis or other protocols well known in the art.

Peptides having the same amino acid sequence as those isolated above may be synthesized by solid-phase peptide synthesis using procedures similar to those described by
35 Merrifield, 1963, *J. Am. Chem. Soc.*, 85:2149. During synthesis, N- α -protected amino acids having protected side chains are added stepwise to a growing polypeptide chain

linked by its C-terminal and to an insoluble polymeric support i.e., polystyrene beads. The peptides are synthesized by linking an amino group of an N- α -deprotected amino acid to an α -carboxy group of an N- α -protected amino acid that has been activated by reacting it with a reagent such as dicyclohexylcarbodiimide. The attachment of a free amino group to the activated carboxyl leads to peptide bond formation. The most commonly used N- α -protecting groups include Boc which is acid labile and Fmoc which is base labile.

Briefly, the C-terminal N- α -protected amino acid is first attached to the polystyrene beads. The N- α -protecting group is then removed. The deprotected α -amino group is coupled to the activated α -carboxylate group of the next N- α -protected amino acid. The process is repeated until the desired peptide is synthesized. The resulting peptides are then cleaved from the insoluble polymer support and the amino acid side chains deprotected. Longer peptides can be derived by condensation of protected peptide fragments. Details of appropriate chemistries, resins, protecting groups, protected amino acids and reagents are well known in the art and so are not discussed in detail herein (See, Atherton, et al., 1989, *Solid Phase Peptide Synthesis: A Practical Approach*, IRL Press, and Bodanszky, 1993, *Peptide Chemistry, A Practical Textbook*, 2nd Ed., Springer-Verlag).

Purification of the resulting peptides is accomplished using conventional procedures, such as preparative HPLC using gel permeation, partition and/or ion exchange chromatography. The choice of appropriate matrices and buffers are well known in the art and so are not described in detail herein.

4.1.5. Exogenous Antigenic Molecules

Antigens or antigenic portions thereof can be selected for use as antigenic molecules, for complexing to hsps, from among those known in the art or determined by immunoassay to be able to bind to antibody or MHC molecules (antigenicity) or generate immune response (immunogenicity). To determine

immunogenicity or antigenicity by detecting binding to antibody, various immunoassays known in the art can be used, including but not limited to competitive and non-competitive assay systems using techniques such as radioimmunoassays, 5 ELISA (enzyme linked immunosorbent assay), "sandwich" immunoassays, immunoradiometric assays, gel diffusion precipitin reactions, immunodiffusion assays, *in vivo* immunoassays (using colloidal gold, enzyme or radioisotope labels, for example), western blots, immunoprecipitation 10 reactions, agglutination assays (e.g., gel agglutination assays, hemagglutination assays), complement fixation assays, immunofluorescence assays, protein A assays, and immunoelectrophoresis assays, etc. In one embodiment, antibody binding is detected by detecting a label on the 15 primary antibody. In another embodiment, the primary antibody is detected by detecting binding of a secondary antibody or reagent to the primary antibody. In a further embodiment, the secondary antibody is labelled. Many means are known in the art for detecting binding in an immunoassay 20 and are envisioned for use. In one embodiment for detecting immunogenicity, T cell-mediated responses can be assayed by standard methods, e.g., *in vitro* cytotoxicity assays or *in vivo* delayed-type hypersensitivity assays.

Potentially useful antigens or derivatives thereof for 25 use as antigenic molecules can also be identified by various criteria, such as the antigen's involvement in neutralization of a pathogen's infectivity (wherein it is desired to treat or prevent infection by such a pathogen) (Norrby, 1985, *Summary*, in *Vaccines 85*, Lerner, et al. (eds.), Cold Spring 30 Harbor Laboratory, Cold Spring Harbor, New York, pp. 388-389), type or group specificity, recognition by patients' antisera or immune cells, and/or the demonstration of protective effects of antisera or immune cells specific for the antigen. In addition, where it is desired to treat or 35 prevent a disease caused by pathogen, the antigen's encoded epitope should preferably display a small or no degree of

antigenic variation in time or amongst different isolates of the same pathogen.

Preferably, where it is desired to treat or prevent cancer, known tumor-specific antigens or fragments or derivatives thereof are used. For example, such tumor specific or tumor-associated antigens include but are not limited to KS 1/4 pan-carcinoma antigen (Perez and Walker, 1990, *J. Immunol.* 142:3662-3667; Bumal, 1988, *Hybridoma* 7(4):407-415); ovarian carcinoma antigen (CA125) (Yu, et al., 1991, *Cancer Res.* 51(2):468-475); prostatic acid phosphate (Tailer, et al., 1990, *Nucl. Acids Res.* 18(16):4928); prostate specific antigen (Henttu and Vihko, 1989, *Biochem. Biophys. Res. Comm.* 160(2):903-910; Israeli, et al., 1993, *Cancer Res.* 53:227-230); melanoma-associated antigen p97 (Estin, et al., 1989, *J. Natl. Cancer Inst.* 81(6):445-446); melanoma antigen gp75 (Vijayasardahl, et al., 1990, *J. Exp. Med.* 171(4):1375-1380); high molecular weight melanoma antigen (Natali, et al., 1987, *Cancer* 59:55-63) and prostate specific membrane antigen.

20 In a specific embodiment, an antigen or fragment or derivative thereof specific to a certain tumor is selected for complexing to hsp and subsequent administration to a patient having that tumor.

Preferably, where it is desired to treat or prevent viral diseases, molecules comprising epitopes of known viruses are used. For example, such antigenic epitopes may be prepared from viruses including, but not limited to, hepatitis type A, hepatitis type B, hepatitis type C, influenza, varicella, adenovirus, herpes simplex type I (HSV-30 I), herpes simplex type II (HSV-II), rinderpest, rhinovirus, echovirus, rotavirus, respiratory syncytial virus, papilloma virus, papova virus, cytomegalovirus, echinovirus, arbovirus, huntavirus, coxsackie virus, mumps virus, measles virus, rubella virus, polio virus, human immunodeficiency virus type 35 I (HIV-I), and human immunodeficiency virus type II (HIV-II).

Preferably, where it is desired to treat or prevent bacterial infections, molecules comprising epitopes of known

bacteria are used. For example, such antigenic epitopes may be prepared from bacteria including, but not limited to, mycobacteria rickettsia, mycoplasma, neisseria and legionella.

5 Preferably, where it is desired to treat or prevent protozoal infections, molecules comprising epitopes of known protozoa are used. For example, such antigenic epitopes may be prepared from protozoa including, but not limited to, leishmania, kokzidioa, and trypanosoma.

10 Preferably, where it is desired to treat or prevent parasitic infections, molecules comprising epitopes of known parasites are used. For example, such antigenic epitopes may be from parasites including, but not limited to, chlamydia and rickettsia.

15

4.1.6. In Vitro Production of Stress Protein-Antigenic Molecule Complexes

In an embodiment in which complexes of hsps and the peptides with which they are endogenously associated *in vivo* are not employed, complexes of hsps to antigenic molecules
20 are produced *in vitro*. As will be appreciated by those skilled in the art, the peptides either isolated by the aforementioned procedures or chemically synthesized or recombinantly produced may be reconstituted with a variety of
25 purified natural or recombinant stress proteins *in vitro* to generate immunogenic non-covalent stress protein-antigenic molecule complexes. Alternatively, exogenous antigens or antigenic/immunogenic fragments or derivatives thereof can be noncovalently complexed to stress proteins for use in the
30 immunotherapeutic or prophylactic vaccines of the invention. A preferred, exemplary protocol for noncovalently complexing a stress protein and an antigenic molecule *in vitro* is discussed below.

Prior to complexing, the hsps are pretreated with ATP or
35 low pH to remove any peptides that may be associated with the hsp of interest. When the ATP procedure is used, excess ATP is removed from the preparation by the addition of apyranase.

as described by Levy, et al., 1991, *Cell* 67:265-274. When the low pH procedure is used, the buffer is readjusted to neutral pH by the addition of pH modifying reagents.

The antigenic molecules (1 μ g) and the pretreated hsp 5 (9 μ g) are admixed to give an approximately 5 antigenic molecule: 1 stress protein molar ratio. Then, the mixture is incubated for 15 minutes to 3 hours at 4° to 45°C in a suitable binding buffer such as one containing 20mM sodium phosphate, pH 7.2, 350mM NaCl, 3mM MgCl₂, and 1mM phenyl methyl sulfonyl fluoride (PMSF). The preparations are centrifuged through a Centricon 10 assembly (Millipore) to remove any unbound peptide. The association of the peptides with the stress proteins can be assayed by SDS-PAGE. This is the preferred method for *in vitro* complexing of peptides isolated from MHC-peptide complexes of peptides disassociated from endogenous hsp-peptide complexes.

In an alternative embodiment of the invention, preferred for producing complexes of hsp70 to exogenous antigenic molecules such as proteins, 5-10 micrograms of purified hsp 20 is incubated with equimolar quantities of the antigenic molecule in 20mM sodium phosphate buffer pH 7.5, 0.5M NaCl, 3mM MgCl₂, and 1mM ADP in a volume of 100 microliter at 37°C for 1 hr. This incubation mixture is further diluted to 1ml in phosphate-buffered saline.

25 In an alternative embodiment of the invention, preferred for producing complexes of gp96 or hsp90 to peptides, 5-10 micrograms of purified gp96 or hsp90 is incubated with equimolar or excess quantities of the antigenic peptide in a suitable buffer such as one containing 20mM sodium phosphate 30 buffer pH 7.5, 0.5M NaCl, 3mM MgCl₂ at 60-65°C for 5-20 min. This incubation mixture is allowed to cool to room temperature and centrifuged one or more times if necessary, through a Centricon 10 assembly (Millipore) to remove any unbound peptide.

35 Following complexing, the immunogenic stress protein-antigenic molecule complexes can optionally be assayed *in vitro* using for example the mixed lymphocyte target cell

assay (MLTC) described below. Once immunogenic complexes have been isolated they can be optionally characterized further in animal models using the preferred administration protocols and excipients discussed below.

5

4.1.7. Determination of Immunogenicity of Stress Protein-Peptide Complexes

The purified stress protein-antigenic molecule complexes can be assayed for immunogenicity using the MLTC assay well known in the art.

10

By way of example but not limitation, the following procedure can be used. Briefly, mice are injected, using any convenient route of administration with the candidate stress protein-antigenic molecule complexes. Other mice are injected with either other stress protein peptide complexes or whole infected cells which act as positive controls for the assay. The mice are injected twice, 7-10 days apart. Ten days after the last immunization, the spleens are removed and the lymphocytes released. The released lymphocytes may be restimulated subsequently in vitro by the addition of dead cells that expressed the complex of interest.

15

For example, 8×10^6 immune spleen cells may be stimulated with 4×10^4 mitomycin C treated or γ -irradiated (5-10,000 rads) infected cells (or cells transfected with an appropriate gene, as the case may be) in 3ml RPMI medium containing 10% fetal calf serum. In certain cases 33% secondary mixed lymphocyte culture supernatant may be included in the culture medium as a source of T cell growth factors (See, Glasebrook, et al., 1980, *J. Exp. Med.* 151:876). To test the primary cytotoxic T cell response after immunization, spleen cells may be cultured without stimulation. In some experiments spleen cells of the immunized mice may also be restimulated with antigenically distinct cells, to determine the specificity of the cytotoxic T cell response.

25

30

35

Six days later the cultures are tested for cytotoxicity in a 4 hour ^{51}Cr -release assay (See, Palladino, et al., 1987, *Cancer-Res.* 47:5074-5079 and Blachere, et al., 1993, *J.*

Immunotherapy 14:352-356). In this assay, the mixed lymphocyte culture is added to a target cell suspension to give different effector:target (E:T) ratios (usually 1:1 to 40:1). The target cells are prelabelled by incubating 1×10^6 target cells in culture medium containing 20 mCi $^{51}\text{Cr}/\text{ml}$ for one hour at 37°C . The cells are washed three times following labeling. Each assay point (E:T ratio) is performed in triplicate and the appropriate controls incorporated to measure spontaneous ^{51}Cr release (no lymphocytes added to assay) and 100% release (cells lysed with detergent). After incubating the cell mixtures for 4 hours, the cells are pelleted by centrifugation at 200g for 5 minutes. The amount of ^{51}Cr released into the supernatant is measured by a gamma counter. The percent cytotoxicity is measured as cpm in the test sample minus spontaneously released cpm divided by the total detergent released cpm minus spontaneously released cpm.

In order to block the MHC class I cascade a concentrated hybridoma supernatant derived from K-44 hybridoma cells (an anti-MHC class I hybridoma) is added to the test samples to a final concentration of 12.5%.

4.1.8. Dosage Regimens

It was established in experimental tumor models (Blachere et al., 1993, *J. Immunotherapy* 14:352-356) that the lowest dose of hsp noncovalently bound to peptide complexes which produced tumor regression in mice was between 10 and 25 microgram/mouse weighing 20-25g which is equal to $25\mu\text{g}/25\text{g} = 1\text{mg}/\text{kg}$. Prior art methods extrapolate to human dosages based on body weight and surface area. For example, prior art methods of extrapolating human dosage based on body weight can be carried out as follows: since the conversion factor for converting the mouse dosage to human dosage is Dose Human per kg = Dose Mouse per kg $\times 12$ (See Freireich, E.J., et al., 1966, *Cancer Chemotherap. Rep.* 50:219-244), the effective dose of hsp-peptide complexes in humans weighing 70kg should be $1\text{mg}/\text{kg} \div 12 \times 70$, i.e., about 6mg (5.8mg).

Drug doses are also given in milligrams per square meter of body surface area because this method rather than body weight achieves a good correlation to certain metabolic and excretory functions (Shirkey, H.C., 1965, *JAMA* 193:443).

5 Moreover, body surface area can be used as a common denominator for drug dosage in adults and children as well as in different animal species as indicated below in Table 1 (Freireich, E.J., et al., 1966, *Cancer Chemotherap. Rep.* 50:219-244).

10

15

20

25

30

35

TABLE 1

**REPRESENTATIVE SURFACE AREA TO WEIGHT
RATIOS (km) FOR VARIOUS SPECIES¹**

5	Species	Body Weight (kg)	Surface Area (Sq m)	km Factor
	Mouse	0.02	0.0066	3.0
	Rat	0.15	0.025	5.9
10	Monkey	3.0	0.24	12
	Dog	8.0	0.40	20
	Human, Child	20	0.80	25
	Adult	60	1.6	37

15 Example: To express a mg/kg dose in any given species as the equivalent mg/sq m dose, multiply the dose by the appropriate km factor. In an adult human, 100 mg/kg is equivalent to 100 mg/kg x 37 kg/sq m = 3700 mg/sq m.

20

¹ Freireich, et al., 1966, Cancer Chemotherap. Rep. 50: 219-244.

25 In contrast to both of the above-described prior art methods of determining dosage levels, dosages of the purified complexes of hsps and antigenic molecules used for administration are preferably much smaller than the dosages estimated by prior art methods. For example, according to a
30 preferred embodiment of the invention, an amount of hsp70- and/or gp96-antigenic molecule complexes is administered subcutaneously that is in the range of about 10 micrograms to about 600 micrograms for a human patient, the more preferred
35 human dosage being the same as used in a 25g mouse, i.e., in the range of 10-100 micrograms. The preferred dosage for hsp-90 peptide complexes in a human patient provided by the present invention is in the range of about 50 to 5,000

micrograms, the more preferred dosage being 100 micrograms. Alternatively, in a specific embodiment, an amount of hsp70- and/or gp96-antigenic molecule complexes is administered intradermally or mucosally that is in the range of about 0.1 5 micrograms to about 60 micrograms for a human patient. In another specific embodiment, the therapeutically effective amount of hsp70- and/or gp96-antigenic molecule complexes is less than 10 micrograms, e.g., in the range of 0.1 to 9 micrograms; the preferred human dosage being substantially 10 equivalent to or smaller than the dosage used in a 25 g mouse, e.g., in the range of 0.5 to 2.0 micrograms. The preferred dosage for hsp90-antigenic molecule complexes for intradermal or mucosal administration to a human patient is in the range of about 5 to 500 micrograms. In a specific 15 embodiment, the therapeutically effective amount of hsp90-antigenic molecule complexes is less than 50 micrograms, e.g., in the range of 5 to 49 micrograms; the preferred dosage being in the range of 5 to 40 micrograms.

The doses recited above can be administered every other 20 day for a total of 5 injections. In a preferred embodiment, the doses recited above are given once weekly for a period of about 4 to 6 weeks, and the mode of administration is preferably varied with each administration. In a preferred example, each site of administration is varied sequentially. 25 Thus, by way of example and not limitation, the first injection may be given intradermally on the left arm, the second on the right arm, the third on the left belly, the fourth on the right belly, the fifth on the left thigh, the sixth on the right thigh, etc. The same site may be repeated 30 after a gap of one or more injections. Also, split injections may be given. Thus, for example, half the dose may be given in one site and the other half in another site on the same day.

After 4-6 weeks, further injections are preferably given 35 at two-week intervals over a period of time of one month. Later injections may be given monthly. The pace of later injections may be modified, depending upon the patient's

clinical progress and responsiveness to the immunotherapy. Alternatively, the mode of administration is sequentially varied, e.g., weekly administrations are given in sequence intradermally or mucosally.

5 The above regimens for administration of hsp-antigenic molecule complexes may occur before, during or after administration of the hsp-antigenic molecule complex-sensitized APC. For example, the mode of therapy can be sequentially varied, e.g., hsp-antigenic molecule complexes
10 may be administered at one time and hsp-antigenic molecule-sensitized APC another time. Alternatively, hsp-antigenic molecule complexes may be administered concurrently with hsp-antigenic molecule-sensitized APC. Preferably, the APC and complexes are administered to the patient within 1 week of
15 each other.

The invention is illustrated by non-limiting examples in Sections 5 and 6.

4.2. Combination With Adoptive Immunotherapy

20 Adoptive immunotherapy refers to a therapeutic approach for treating cancer or infectious diseases in which immune cells are administered to a host with the aim that the cells mediate either directly or indirectly specific immunity to tumor cells and/or antigenic components or regression of the
25 tumor or treatment of infectious diseases, as the case may be. (See U.S. Patent Application Serial No. 08/527,546, filed September 13, 1995, which is incorporated by reference herein in its entirety.) In accordance with the methods described herein, APC are sensitized with hsps noncovalently
30 complexed with antigenic (or immunogenic) molecules and used in adoptive immunotherapy.

According to the invention, therapy by administration of hsp-peptide complexes, using any desired route of administration, is combined with adoptive immunotherapy using
35 APC sensitized with hsp-antigenic molecule complexes. As described in Section 4 herein, the hsp-peptide complex-sensitized APC can be administered concurrently with hsp-

peptide complexes, or before or after administration of hsp-peptide complexes. Furthermore, the mode of administration can be varied, including but not limited to, e.g., subcutaneously, intravenously, intraperitoneally, 5 intramuscularly, intradermally or mucosally.

4.2.1. Obtaining Macrophages and Antigen-Presenting Cells

The antigen-presenting cells, including but not limited 10 to macrophages, dendritic cells and B-cells, are preferably obtained by production *in vitro* from stem and progenitor cells from human peripheral blood or bone marrow as described by Inaba, K., et al., 1992, J. Exp. Med. 176:1693-1702.

APC can be obtained by any of various methods known in 15 the art. In a preferred aspect human macrophages are used, obtained from human blood cells. By way of example but not limitation, macrophages can be obtained as follows:

Mononuclear cells are isolated from peripheral blood of a patient (preferably the patient to be treated), by Ficoll- 20 Hypaque gradient centrifugation and are seeded on tissue culture dishes which are pre-coated with the patient's own serum or with other AB+ human serum. The cells are incubated at 37°C for 1 hr, then non-adherent cells are removed by pipetting. To the adherent cells left in the dish, is added 25 cold (4°C) 1 mM EDTA in phosphate-buffered saline and the dishes are left at room temperature for 15 minutes. The cells are harvested, washed with RPMI buffer and suspended in RPMI buffer. Increased numbers of macrophages may be obtained by incubating at 37°C with macrophage-colony 30 stimulating factor (M-CSF); increased numbers of dendritic cells may be obtained by incubating with granulocyte-macrophage-colony stimulating factor (GM-CSF) as described in detail by Inaba, K., et al., 1992, J. Exp. Med. 176:1693-1702.

35

4.2.2. Sensitization of Macrophages and Antigen Presenting Cells With Hsp-Peptide Complexes

APC are sensitized with hsp noncovalently bound to antigenic molecules by incubating the cells in vitro with the complexes. The APC are sensitized with complexes of hsps and antigenic molecules preferably by incubating in vitro with the hsp-complex at 37°C for 15 minutes to 24 hours. By way of example but not limitation, 4×10^7 macrophages can be incubated with 10 microgram gp96-peptide complexes per ml or 100 microgram hsp90-peptide complexes per ml at 37°C for 15 minutes-24 hours in 1 ml plain RPMI medium. The cells are washed three times and resuspended in a physiological medium preferably sterile, at a convenient concentration (e.g., 1×10^7 /ml) for injection in a patient. Preferably, the patient into which the sensitized APCs are injected is the patient from which the APC were originally isolated (autologous embodiment).

Optionally, the ability of sensitized APC to stimulate, for example, the antigen-specific, class I-restricted cytotoxic T-lymphocytes (CTL) can be monitored by their ability to stimulate CTLs to release tumor necrosis factor, and by their ability to act as targets of such CTLs.

4.2.3. Reinfusion of Sensitized APC

The hsp-antigenic molecule-sensitized APC are reinfused into the patient systemically, preferably intravenously, by conventional clinical procedures. These activated cells are reinfused, preferentially by systemic administration into the autologous patient. Patients generally receive from about 10^6 to about 10^{12} sensitized macrophages, depending on the condition of the patient. In some regimens, patients may optionally receive in addition a suitable dosage of a biological response modifier including but not limited to the cytokines IFN- α , IFN- γ , IL-2, IL-4, IL-6, TNF or other cytokine growth factor.

4.3. Formulation, Administration & Kits

Kits of the invention are provided that comprise in a first container a pharmaceutical composition comprising a complex consisting essentially of a heat shock protein noncovalently bound to an antigenic molecule and a pharmaceutically acceptable carrier; and in a second container antigen presenting cells. Hsp-antigenic molecule complexes of the invention may be formulated into pharmaceutical preparations for administration to mammals, preferably humans, for treatment or prevention of cancer or infectious diseases. Compositions comprising a compound of the invention formulated in a compatible pharmaceutical carrier may be prepared, packaged, and labelled for treatment of the indicated tumor(s), such as human sarcomas and carcinomas, e.g., fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteogenic sarcoma, chordoma, angiosarcoma, endotheliosarcoma, lymphangiosarcoma, lymphangioendotheliosarcoma, synovioma, mesothelioma, Ewing's tumor, leiomyosarcoma, rhabdomyosarcoma, colon carcinoma, pancreatic cancer, breast cancer, ovarian cancer, prostate cancer, squamous cell carcinoma, basal cell carcinoma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma, papillary adenocarcinomas, cystadenocarcinoma, medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma, hepatoma, bile duct carcinoma, choriocarcinoma, seminoma, embryonal carcinoma, Wilms' tumor, cervical cancer, testicular tumor, lung carcinoma, small cell lung carcinoma, bladder carcinoma, epithelial carcinoma, glioma, astrocytoma, medulloblastoma, craniopharyngioma, ependymoma, pinealoma, hemangioblastoma, acoustic neuroma, oligodendroglioma, meningioma, melanoma, neuroblastoma, retinoblastoma; leukemias, e.g., acute lymphocytic leukemia and acute myelocytic leukemia (myeloblastic, promyelocytic, myelomonocytic, monocytic and erythroleukemia); chronic leukemia (chronic myelocytic (granulocytic) leukemia and chronic lymphocytic leukemia); and polycythemia vera, lymphoma (Hodgkin's disease and non-

Hodgkin's disease), multiple myeloma, Waldenström's macroglobulinemia, and heavy chain disease. Alternatively, it can be labeled for treatment of the appropriate infectious disease. Alternatively, pharmaceutical compositions may be
5 formulated for treatment of appropriate infectious diseases.

Drug solubility and the site of absorption are factors which should be considered when choosing the route of administration of a therapeutic agent. In an embodiment of the invention, hsp-antigenic molecule complexes may be
10 administered using any desired route of administration. Advantages of intradermal or mucosal administration include use of lower doses and rapid absorption, respectively. Advantages of subcutaneous or intramuscular administration include suitability for some insoluble suspensions and oily
15 suspensions, respectively. Mucosal routes of administration include, but are not limited to, oral, rectal and nasal administration. Preparations for mucosal administrations are suitable in various formulations as described below.

If the complex is water-soluble, then it may be
20 formulated in an appropriate buffer, for example, phosphate buffered saline or other physiologically compatible solutions, preferably sterile. Alternatively, if the resulting complex has poor solubility in aqueous solvents, then it may be formulated with a non-ionic surfactant such as
25 Tween, or polyethylene glycol. Thus, the compounds and their physiologically acceptable solvates may be formulated for administration by inhalation or insufflation (either through the mouth or the nose) or oral, buccal, parenteral, or rectal administration or, in the case of tumors, directly injected
30 into a solid tumor.

For oral administration, the pharmaceutical preparation may be in liquid form, for example, solutions, syrups or suspensions, or may be presented as a drug product for reconstitution with water or other suitable vehicle before
35 use. Such liquid preparations may be prepared by conventional means with pharmaceutically acceptable additives such as suspending agents (e.g., sorbitol syrup, cellulose

derivatives or hydrogenated edible fats); emulsifying agents (e.g., lecithin or acacia); non-aqueous vehicles (e.g., almond oil, oily esters, or fractionated vegetable oils); and preservatives (e.g., methyl or propyl-p-hydroxybenzoates or sorbic acid). The pharmaceutical compositions may take the form of, for example, tablets or capsules prepared by conventional means with pharmaceutically acceptable excipients such as binding agents (e.g., pregelatinized maize starch, polyvinyl pyrrolidone or hydroxypropyl methylcellulose); fillers (e.g., lactose, microcrystalline cellulose or calcium hydrogen phosphate); lubricants (e.g., magnesium stearate, talc or silica); disintegrants (e.g., potato starch or sodium starch glycolate); or wetting agents (e.g., sodium lauryl sulphate). The tablets may be coated by methods well-known in the art.

Preparations for oral administration may be suitably formulated to give controlled release of the active compound.

For buccal administration, the compositions may take the form of tablets or lozenges formulated in conventional manner.

The compounds may be formulated for parenteral administration by injection, e.g., by bolus injection or continuous infusion. Formulations for injection may be presented in unit dosage form, e.g., in ampoules or in multi-dose containers, with an added preservative. The compositions may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents. Alternatively, the active ingredient may be in powder form for constitution with a suitable vehicle, e.g., sterile pyrogen-free water, before use.

The compounds may also be formulated in rectal compositions such as suppositories or retention enemas, e.g., containing conventional suppository bases such as cocoa butter or other glycerides.

In addition to the formulations described previously, the compounds may also be formulated as a depot preparation.

Such long acting formulations may be administered by implantation (for example, subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the compounds may be formulated with suitable polymeric or hydrophobic materials (for example, as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt. Liposomes and emulsions are well known examples of delivery vehicles or carriers for hydrophilic drugs.

- 10 For administration by inhalation, the compounds for use according to the present invention are conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebulizer, with the use of a suitable propellant, e.g., dichlorodifluoromethane, trichlorofluoromethane, 15 dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and cartridges of, e.g., gelatin for use in an inhaler or insufflator may be formulated containing a 20 powder mix of the compound and a suitable powder base such as lactose or starch.

The compositions may, if desired, be presented in a pack or dispenser device which may contain one or more unit dosage forms containing the active ingredient. The pack may for 25 example comprise metal or plastic foil, such as a blister pack. The pack or dispenser device may be accompanied by instructions for administration.

- The invention also provides kits for carrying out the therapeutic regimens of the invention. Such kits comprise in 30 a first one or more containers therapeutically or prophylactically effective amounts of the hsp-antigenic molecule complexes, preferably purified, in pharmaceutically acceptable form; and in a second container the sensitized APC of the invention, preferably purified. The hsp-antigenic 35 molecule complex in a vial of a kit of the invention may be in the form of a pharmaceutically acceptable solution, e.g., in combination with sterile saline, dextrose solution, or

buffered solution, or other pharmaceutically acceptable sterile fluid. Alternatively, the complex may be lyophilized or desiccated; in this instance, the kit optionally further comprises in a container a pharmaceutically acceptable
5 solution (e.g., saline, dextrose solution, etc.), preferably sterile, to reconstitute the complex to form a solution for injection purposes.

In another embodiment, a kit of the invention further comprises a needle or syringe, preferably packaged in sterile
10 form, for injecting the complex, and/or a packaged alcohol pad. Instructions are optionally included for administration of hsp-antigenic molecule complexes by a clinician or by the patient.

15 **4.4. Target Infectious Diseases**

Infectious diseases that can be treated or prevented by the methods of the present invention are caused by infectious agents including, but not limited to, viruses, bacteria, fungi protozoa and parasites.

20 Viral diseases that can be treated or prevented by the methods of the present invention include, but are not limited to, those caused by hepatitis type A, hepatitis type B, hepatitis type C, influenza, varicella, adenovirus, herpes simplex type I (HSV-I), herpes simplex type II (HSV-II),
25 rinderpest, rhinovirus, echovirus, rotavirus, respiratory syncytial virus, papilloma virus, papova virus, cytomegalovirus, echinovirus, arbovirus, huntavirus, coxsackie virus, mumps virus, measles virus, rubella virus, polio virus, human immunodeficiency virus type I (HIV-I), and
30 human immunodeficiency virus type II (HIV-II).

Bacterial diseases that can be treated or prevented by the methods of the present invention are caused by bacteria including, but not limited to, mycobacteria rickettsia, mycoplasma, neisseria and legionella.

35 Protozoal diseases that can be treated or prevented by the methods of the present invention are caused by protozoa

including, but not limited to, leishmania, kokzidioa, and trypanosoma.

Parasitic diseases that can be treated or prevented by the methods of the present invention are caused by parasites including, but not limited to, chlamydia and rickettsia.

4.5. Target Cancers

Cancers that can be treated or prevented by the methods of the present invention include, but are not limited to

10 human sarcomas and carcinomas, e.g., fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteogenic sarcoma, chordoma, angiosarcoma, endotheliosarcoma, lymphangiosarcoma, lymphangioendotheliosarcoma, synovioma, mesothelioma, Ewing's tumor, leiomyosarcoma, rhabdomyosarcoma, colon carcinoma,

15 pancreatic cancer, breast cancer, ovarian cancer, prostate cancer, squamous cell carcinoma, basal cell carcinoma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma, papillary adenocarcinomas, cystadenocarcinoma, medullary carcinoma, bronchogenic

20 carcinoma, renal cell carcinoma, hepatoma, bile duct carcinoma, choriocarcinoma, seminoma, embryonal carcinoma, Wilms' tumor, cervical cancer, testicular tumor, lung carcinoma, small cell lung carcinoma, bladder carcinoma, epithelial carcinoma, glioma, astrocytoma, medulloblastoma,

25 craniopharyngioma, ependymoma, pinealoma, hemangioblastoma, acoustic neuroma, oligodendroglioma, meningioma, melanoma, neuroblastoma, retinoblastoma; leukemias, e.g., acute lymphocytic leukemia and acute myelocytic leukemia (myeloblastic, promyelocytic, myelomonocytic, monocytic and

30 erythroleukemia); chronic leukemia (chronic myelocytic (granulocytic) leukemia and chronic lymphocytic leukemia); and polycythemia vera, lymphoma (Hodgkin's disease and non-Hodgkin's disease), multiple myeloma, Waldenström's macroglobulinemia, and heavy chain disease. Specific

35 examples of such cancers are described in the sections below.

In a specific embodiment the cancer is metastatic. In another specific embodiment, the patient having a cancer is

immunosuppressed by reason of having undergone anti-cancer therapy (e.g., chemotherapy radiation) prior to administration of the hsp-antigenic molecule complexes or administration of the hsp-sensitized APC.

5

4.5.1. Colorectal Cancer Metastatic to the Liver

In 1992, approximately 150,000 Americans were diagnosed with colorectal cancer and more than 60,000 died as a result of colorectal metastases. At the time of their deaths, 80 percent of patients with colorectal cancer have metastatic disease involving the liver, and one-half of these patients have no evidence of other (extrahepatic) metastases. Most metastatic tumors of the liver are from gastrointestinal primaries. Unfortunately, the natural history of metastatic liver lesions carries a grave prognosis and systemic chemotherapy regimens have been unable to induce significant response rates or alter length of survival (Drebin, J.A., et al., in *Current Therapy In Oncology*, ed. J.E. Niederhuber, B.C. Decker, Mosby, 1993, p.426).

20

Colorectal cancer initially spreads to regional lymph nodes and then through the portal venous circulation to the liver, which represents the most common visceral site of metastasis. The symptoms that lead patients with colorectal cancer to seek medical care vary with the anatomical location of the lesion. For example, lesions in the ascending colon frequently ulcerate, which leads to chronic blood loss in the stool.

25

Radical resection offers the greatest potential for cure in patients with invasive colorectal cancer. Before surgery, the CEA titer is determined. Radiation therapy and chemotherapy are used in patients with advanced colorectal cancer. Results with chemotherapeutic agents (e.g., 5-fluorouracil) are mixed and fewer than 25 percent of patients experience a greater than 50 percent reduction in tumor mass (Richards, 2d., F., et al., 1986, *J. Clin. Oncol.* 4:565).

35

Patients with widespread metastases have limited survival and systemic chemotherapy has little impact in this group of patients. In addition, systemically administered chemotherapy is often limited by the severity of toxicities 5 associated with the various agents, such as severe diarrhea, mucositis and/or myelosuppression. Other techniques, including hepatic radiation, systemic chemotherapy, hepatic arterial ligation, tumor embolization and immunotherapy have all been explored, but, for the most part, have proven 10 ineffectual in prolonging patient survival.

In a specific embodiment, the present invention provides compositions and methods for enhancing tumor specific immunity in individuals suffering from colorectal cancer metastasized to the liver, in order to inhibit the 15 progression of the neoplastic disease. Preferred methods of treating these neoplastic diseases comprise administering a composition of autologous hsp noncovalently bound to peptide complexes, which elicits tumor-specific immunity against the tumor cells. Most specifically, the use of a composition of 20 the invention, comprising gp96, can result in nearly complete inhibition of liver cancer growth in cancer patients, without inducing toxicity and thus providing a dramatic therapeutic effect.

Accordingly, as an example of the method of the 25 invention, gp96-antigenic molecule complexes are administered to a patient diagnosed with colorectal cancer, with or without liver metastasis, via one of many different routes of administration, the preferred route being intradermally at different anatomical sites, e.g., left arm, right arm, left 30 belly, right belly, left thigh, right thigh, etc. The site of injection is varied for each weekly injection as described in Sections 5 and 6. Exemplary primary and metastatic cancers that can be prevented or treated according to the methods of the invention are described in detail in the 35 sections which follow and by way of example, *infra*.

4.5.2. Hepatocellular Carcinoma

Hepatocellular carcinoma is generally a disease of the elderly in the United States. Although many factors may lead to hepatocellular carcinoma, the disease is usually limited to those persons with preexisting liver disease.

Approximately 60 to 80 percent of patients in the United States with hepatocellular carcinoma have a cirrhotic liver and about four percent of individuals with a cirrhotic liver eventually develop hepatocellular carcinoma (Niederhuber, J.E., (ed.), 1993, *Current Therapy in Oncology*, B.C. Decker, Mosby). The risk is highest in patients whose liver disease is caused by inherited hemochromatosis or hepatic B viral infection (Bradbear, R.A., et al., 1985, *J. Natl. Cancer Inst.* 75:81; Beasley, R.P., et al., 1981, *Lancet* 2:1129).

Other causes of cirrhosis that can lead to hepatocellular carcinoma include alcohol abuse and hepatic fibrosis caused by chronic administration of methotrexate. The most frequent symptoms of hepatocellular carcinoma are the development of a painful mass in the right upper quadrant or epigastrium, accompanied by weight loss. In patients with cirrhosis, the development of hepatocellular carcinoma is preceded by ascites, portal hypertension and relatively abrupt clinical deterioration. In most cases, abnormal values in standard liver function tests such as serum aminotransferase and alkaline phosphatase are observed.

CT scans of the liver are used to determine the anatomic distribution of hepatocellular carcinoma and also provide orientation for percutaneous needle biopsy. Approximately 70 percent of patients with hepatocellular carcinoma have an elevated serum alpha-fetoprotein concentration (McIntire, K.R., et al., 1975, *Cancer Res.* 35:991) and its concentration correlates with the extent of the disease.

Radical resection offers the only hope for cure in patients with hepatocellular carcinoma. Such operative procedures are associated with five-year survival rates of 12 to 30 percent. Liver transplantation may improve survival of some younger individuals. However, most patients are not

surgical candidates because of extensive cirrhosis multifocal tumor pattern or scarcity of compatible donor organs. Chemotherapeutic agents have been administered either by intravenous route or through an intrahepatic arterial catheter. Such therapy has sometimes been combined with irradiation to the liver. Reductions in the size of measurable tumors of 50% or more have been reported in some patients treated with either systemic doxorubicin or 5-fluorouracil. However, chemotherapy often induces immunosuppression and rarely causes the tumor to disappear completely and the duration of response is short. The prognosis for patients with hepatocellular carcinoma is negatively correlated with cirrhosis and metastases to the lungs or bone. Median survival for patients is only four to six months. In another specific embodiment, the present invention provides compositions and methods for enhancing specific immunity in individuals suffering from hepatocellular carcinoma in order to inhibit the progression of the neoplastic disease and ultimately irradiate all preneoplastic and neoplastic cells.

4.5.3. Breast Cancer

Another specific aspect of the invention relates to the treatment of breast cancer. The American Cancer Society estimated that in 1992 180,000 American women were diagnosed with breast cancer and 46,000 succumbed to the disease (Niederhuber, J.E.ed. *Current Therapy in Oncology* B.C. Decker, Mosby, 1993). This makes breast cancer the second major cause of cancer death in women, ranking just behind lung cancer. A disturbing fact is the observation that breast cancer has been increasing at a rate of 3 percent per year since 1980 (Niederhuber, J.E., ed. Current Therapy in Oncology, B.C. Decker, Mosby, (1993)). The treatment of breast cancer presently involves surgery, radiation, hormonal therapy and/or chemotherapy. Consideration of two breast cancer characteristics, hormone receptors and disease extent, has governed how hormonal therapies and standard-dose

chemotherapy are sequenced to improve survival and maintain or improve quality of life. A wide range of multidrug regimens have been used as adjuvant therapy in breast cancer patients, including, but not limited to combinations of 2
5 cyclophosphamide, doxorubicin, vincristine methotrexate, 5-fluorouracil and/or leucovorin. In a specific embodiment, the present invention provides hsp compositions and methods for enhancing specific immunity to preneoplastic and neoplastic mammary cells in women. The present invention
10 also provides compositions and methods for preventing the development of neoplastic cells in women at enhanced risk for breast cancer, and for inhibiting cancer cell proliferation and metastasis. These compositions can be applied alone or in combination with each other or with biological response
15 modifiers.

4.6. Autologous Embodiment

The specific immunogenicity of hsps derives not from hsps per se, but from the peptides bound to them. In a
20 preferred embodiment of the invention directed to the use of autologous complexes of hsp-peptides as cancer vaccines, two of the most intractable hurdles to cancer immunotherapy are circumvented. First is the possibility that human cancers, like cancers of experimental animals, are antigenically
25 distinct. In an embodiment of the present invention, hsps chaperone antigenic peptides of the cancer cells from which they are derived and circumvent this hurdle. Second, most current approaches to cancer immunotherapy focus on determining the CTL-recognized epitopes of cancer cell lines.
30 This approach requires the availability of cell lines and CTLs against cancers. These reagents are unavailable for an overwhelming proportion of human cancers. In an embodiment of the present invention directed to the use of autologous complexes of hsp-peptides, cancer immunotherapy does not
35 depend on the availability of cell lines or CTLs nor does it require definition of the antigenic epitopes of cancer cells.

These advantages make autologous hsps noncovalently bound to peptide complexes attractive immunogens against cancer.

4.7. Prevention and Treatment of Primary and Metastatic Neoplastic Diseases

5 There are many reasons why immunotherapy as provided by the present invention is desired for use in cancer patients. First, if cancer patients are immunosuppressed, surgery with anesthesia and subsequent chemotherapy may worsen the
10 immunosuppression. With appropriate immunotherapy in the preoperative period, this immunosuppression may be prevented or reversed. This could lead to fewer infectious complications and to accelerated wound healing. Second, tumor bulk is minimal following surgery and immunotherapy is
15 most likely to be effective in this situation. A third reason is the possibility that tumor cells are shed into the circulation at surgery and effective immunotherapy applied at this time can eliminate these cells.

The preventive and therapeutic methods of the invention
20 are directed at enhancing the immunocompetence of the cancer patient either before surgery, at or after surgery, and to induce tumor-specific immunity to cancer cells, with the objective being inhibition of cancer, and with the ultimate clinical objective being total cancer regression and
25 eradication.

4.8. Monitoring of Effects During Cancer Prevention and Immunotherapy with Hsp-peptide Complexes in Combination with Adoptive Immunotherapy

The effect of immunotherapy with hsp-antigenic molecule
30 complexes on development and progression of neoplastic diseases can be monitored by any methods known to one skilled in the art, including but not limited to measuring: a) delayed hypersensitivity as an assessment of cellular
immunity; b) activity of cytolytic T-lymphocytes *in vitro*; c)
35 levels of tumor specific antigens, e.g., carcinoembryonic (CEA) antigens; d) changes in the morphology of tumors using techniques such as a computed tomographic (CT) scan; and e)

changes in levels of putative biomarkers of risk for a particular cancer in individuals at high risk, and f) changes in the morphology of tumors using a sonogram.

5 4.8.1. Delayed Hypersensitivity Skin Test

Delayed hypersensitivity skin tests are of great value in the overall immunocompetence and cellular immunity to an antigen. Inability to react to a battery of common skin antigens is termed anergy (Sato, T., et al, 1995, Clin. Immunol. Pathol. 74:35-43).

Proper technique of skin testing requires that the antigens be stored sterile at 4°C, protected from light and reconstituted shortly before use. A 25- or 27-gauge needle ensures intradermal, rather than subcutaneous, administration of antigen. Twenty-four and 48 hours after intradermal administration of the antigen, the largest dimensions of both erythema and induration are measured with a ruler. Hypoactivity to any given antigen or group of antigens is confirmed by testing with higher concentrations of antigen or, in ambiguous circumstances, by a repeat test with an intermediate test.

 4.8.2. Activity of Cytolytic T-lymphocytes In Vitro

25 8x10⁶ peripheral blood derived T lymphocytes isolated by the Ficoll-Hypaque centrifugation gradient technique, are restimulated with 4x10⁴ mitomycin C treated tumor cells in 3ml RPMI medium containing 10% fetal calf serum. In some experiments, 33% secondary mixed lymphocyte culture supernatant or IL-2, is included in the culture medium as a source of T cell growth factors.

30 In order to measure the primary response of cytolytic T-lymphocytes after immunization, T cells are cultured without the stimulator tumor cells. In other experiments, T cells are restimulated with antigenically distinct cells. After six days, the cultures are tested for cytotoxicity in a 4 hour ⁵¹Cr-release assay. The spontaneous ⁵¹Cr-release of the

targets should reach a level less than 20%. For the anti-MHC class I blocking activity, a tenfold concentrated supernatant of W6/32 hybridoma is added to the test at a final concentration of 12.5% (Heike M., et al., *J. Immunotherapy* 5 15:165-174).

4.8.3. Levels of Tumor Specific Antigens

Although it may not be possible to detect unique tumor antigens on all tumors, many tumors display antigens that
10 distinguish them from normal cells. The monoclonal antibody reagents have permitted the isolation and biochemical characterization of the antigens and have been invaluable diagnostically for distinction of transformed from
nontransformed cells and for definition of the cell lineage
15 of transformed cells. The best-characterized human tumor-associated antigens are the oncofetal antigens. These antigens are expressed during embryogenesis, but are absent or very difficult to detect in normal adult tissue. The prototype antigen is carcinoembryonic antigen (CEA), a
20 glycoprotein found on fetal gut and human colon cancer cells, but not on normal adult colon cells. Since CEA is shed from colon carcinoma cells and found in the serum, it was originally thought that the presence of this antigen in the serum could be used to screen patients for colon cancer.
25 However, patients with other tumors, such as pancreatic and breast cancer, also have elevated serum levels of CEA. Therefore, monitoring the fall and rise of CEA levels in cancer patients undergoing therapy has proven useful for predicting tumor progression and responses to treatment.
30 Several other oncofetal antigens have been useful for diagnosing and monitoring human tumors, e.g., alpha-fetoprotein, an alpha-globulin normally secreted by fetal liver and yolk sac cells, is found in the serum of patients with liver and germinal cell tumors and can be used as a
35 marker of disease status.

4.8.4. Computed Tomographic (CT) Scan

CT remains the choice of techniques for the accurate staging of cancers. CT has proved more sensitive and specific than any other imaging techniques for the detection of metastases.

4.8.5. Measurement of Putative Biomarkers

The levels of a putative biomarker for risk of a specific cancer are measured to monitor the effect of hsp noncovalently bound to peptide complexes. For example, in individuals at enhanced risk for prostate cancer, serum prostate-specific antigen (PSA) is measured by the procedure described by Brawer, M.K., et. al., 1992, J. Urol. 147:841-845, and Catalona, W.J., et al., 1993, JAMA 270:948-958; or in individuals at risk for colorectal cancer, CEA is measured as described above in Section 4.9.3; and in individuals at enhanced risk for breast cancer, 16- α -hydroxylation of estradiol is measured by the procedure described by Schneider, J. et al., 1982, Proc. Natl. Acad. Sci. USA 79:3047-3051.

4.8.6. Sonogram

A sonogram remains an alternative choice of technique for the accurate staging of cancers.

5. EXAMPLES: ADOPTIVE TRANSFER OF SENSITIZED MACROPHAGES, IN COMBINATION WITH ADMINISTRATION OF HSP-PEPTIDE COMPLEXES

Autologous human macrophages are sensitized with autologous human gp96 noncovalently bound to an antigenic/immunogenic molecule. The sensitized macrophages are administered to the human patient at approximately the same time as, or before, or after the administration of the gp96-antigenic molecule complex.

5.1. Materials and Methods

Macrophages are obtained as follows: mononuclear cells are isolated from peripheral blood of the human patient to be treated, by Ficoll-Hypaque gradient centrifugation and are seeded on tissue culture dishes which are pre-coated with the patient's own serum or with other AB+ human serum. The cells are incubated at 37°C for 1 hour, then non-adherent cells are removed by pipetting. To the adherent cells left in the dish, is added cold (4°C) 1mM EDTA in phosphate-buffered saline and the dishes are left at room temperature for 15 minutes. The cells are harvested, washed with RPMI buffer and suspended in RPMI buffer. Increased numbers of macrophages may be obtained by incubating at 37°C with macrophage-colony stimulating factor (M-CSF); increased numbers of dendritic cells may be obtained by incubating with granulocyte-macrophage-colony stimulating factor (GM-CSF) as described in detail by Inaba, K., et al., 1992, J. Exp. Med. 176:1693-1702.

The macrophages (4×10^7) are then incubated at 37°C for 3 hr in 1ml RPMI containing 50µg gp96-peptide complexes derived from the autologous tumor or from autologous liver, using methods as described in Section 4.1.3. The macrophages are then washed 3 times and resuspended at a concentrate of 1×10^7 /ml in RPMI medium. 200 microliters of this suspension is administered as described in the experimental protocol below.

5.2. TREATMENT OF HEPATOCELLULAR CARCINOMA

Five groups of human patients with hepatocellular carcinoma are injected with autologous macrophages sensitized with hsp-peptide complexes derived from their own tumors post surgery. Treatment with hsp-peptide complexes is started any time after surgery. However, if the patient has received chemotherapy, sensitized macrophages alone or in combination with an hsp-peptide complexes are usually administered after an interval of four weeks or more so as to allow the immune

system to recover. The immunocompetence of the patient is tested by procedures described in sections 4.7 above.

The preferred therapeutic regimen includes weekly injections of the sensitized macrophages in combination with an hsp-peptide complex dissolved in saline or other physiologically compatible solution. Sensitized macrophages may be administered at approximately the same time with an hsp-peptide complex or one may be administered prior to administration of the other.

10 The dosage used for hsp70 or gp96 is in the range of 0.1 to 9 micrograms, with the preferred dosage being 0.5-2.0 micrograms. The dosage used for hsp90 is in the range of 5 to 500 micrograms, with the preferred dosage being about 10 micrograms.

15 The site of injection is varied each time, for example, the first injection is given intradermally on the left arm, the second injection intradermally on the right arm, the third injection intradermally on the left abdominal region, the fourth injection intradermally on the right abdominal region, the fifth injection intradermally on the left thigh, the sixth injection intradermally on the right thigh, etc. The same site can be alternatively repeated after a gap of one or more injections. In addition, injections are split and each half of the dose is administered at a different site 25 on the same day.

Overall, the first four to six injections are given at weekly intervals. Subsequently, two injections are given at two-week intervals; followed by a regimen of injections at monthly intervals. The effect of therapy is monitored by measuring: a) delayed hypersensitivity as an assessment of cellular immunity; b) activity of cytolytic T-lymphocytes in vitro; c) levels of tumor specific antigens, e.g., carcinoembryonic (CEA) antigens; d) changes in the morphology of tumors using techniques such as a computed tomographic (CT) scan; and/or e) changes in putative biomarkers of risk 35 for a particular cancer in individuals at high risk.

Depending on the results obtained, as described above in Section 4.10, the therapeutic regimen may be modified to maintain and/or boost the immunological responses of the patient, with the ultimate goal of achieving tumor regression and complete eradication of cancer cells.

6. **EXAMPLE: ADMINISTRATION OF HSP-PEPTIDE COMPLEXES
IN COMBINATION WITH ADOPTIVE IMMUNOTHERAPY
IN THE TREATMENT OF COLORECTAL CANCER**

Hsp-peptide complexes (gp96, hsp70, hsp90 or a combination thereof) are administered in combination with the administration of antigen presenting cells sensitized with an hsp-antigenic molecule complex as adjuvant therapy and as prophylactic adjuvant therapy in patients after complete reduction of colorectal cancer to eliminate undetectable micrometastases and to improve survival.

The therapeutic and prophylactic regimens used in patients suffering from colorectal cancer are the same as those described in Section 5 above for patients recovering with hepatocellular carcinoma. The methods of monitoring of patients under clinical evaluation for prevention and treatment of colorectal cancer is done by procedures described in Section 4.7. Specifically, CEA levels are measured as a useful monitor of tumor regression and/or recurrence (Mayer, R.J., et al., 1978, *Cancer* 42:1428).

The present invention is not to be limited in scope by the specific embodiments described herein. Indeed, various modifications of the invention in addition to those described herein will become apparent to those skilled in the art from the foregoing description and accompanying figures. Such modifications are intended to fall within the scope of the appended claims.

Various publications are cited herein, the disclosures of which are incorporated by reference in their entireties.

WHAT IS CLAIMED IS:

1. A method of eliciting an immune response in an individual comprising:
 - (a) administering to the individual a first composition comprising a first complex consisting essentially of a first heat shock protein noncovalently bound to a first antigenic molecule; and
 - (b) administering to the individual a second composition comprising antigen presenting cells sensitized *in vitro* with a sensitizing amount of a second complex of a second heat shock protein noncovalently bound to a second antigenic molecule; in which said second composition is administered before, concurrently or after administration of the first composition.
2. The method according to claim 1 in which the first heat shock protein is selected from the group consisting of hsp70, hsp90, gp96, and a mixture of two or more of the foregoing.
3. The method according to claim 1 in which the second heat shock protein is selected from the group consisting of hsp70, hsp90, gp96, and a mixture of two or more of the foregoing.
4. The method according to claim 2 in which the second heat shock protein is selected from the group consisting of hsp70, hsp90, gp96, and a mixture of two or more of the foregoing.
5. The method according to claim 1 in which the first heat shock protein is hsp70.
6. The method according to claim 1 in which the first heat shock protein is hsp90.

7. The method according to claim 1 in which the first heat shock protein is gp96.

8. The method according to claim 1 or 2 in which the first heat shock protein and the second heat shock protein are the same.

9. The method according to claim 1 or 2 in which the individual is a human.

10

10. The method according to claim 1 in which the first composition is the same as the second composition.

11. The method according to claim 1 or 2 in which the individual has liver cancer, colon cancer, or breast cancer.

12. The method according to claim 1 or 2, further comprising administering to the individual an effective amount of a biological response modifier selected from the group consisting of interferon- α , interferon- γ , interleukin-2, interleukin-4, interleukin-6, and tumor necrosis factor.

13. The method according to claim 1 or 2 in which the first composition is administered at weekly intervals.

25

14. The method according to claim 1 or 2 in which the second composition is administered at weekly intervals.

15. The method according to claim 1 or 2 in which the first complex is administered intramuscularly, subcutaneously, intraperitoneally, intravenously, intradermally or mucosally.

16. The method according to claim 1 or 2 in which the second composition is administered intravenously.

17. The method according to claim 1 or 2 in which 10^6 to 10^{12} antigen presenting cells are administered.

18. The method according to claim 1 or 2 in which the first and second complexes are the same.

19. The method according to claim 4 in which the individual is a human.

20. A method of treating an individual having cancer comprising:

(a) administering to the individual a first composition comprising a first complex consisting essentially of a first heat shock protein noncovalently bound to a first antigenic molecule; and

(b) administering to the individual a second composition comprising antigen presenting cells sensitized in vitro with a sensitizing amount of a second complex of a second heat shock protein noncovalently bound to a second antigenic molecule; in which said second composition is administered before, concurrently or after administration of the first composition.

21. The method according to claim 20 in which the first heat shock protein is selected from the group consisting of hsp70, hsp90, gp96, and a mixture of two or more of the foregoing.

22. The method according to claim 20 in which the second heat shock protein is selected from the group consisting of hsp70, hsp90, gp96, and a mixture of two or more of the foregoing.

23. The method according to claim 21 in which the second heat shock protein is selected from the group

consisting of hsp70, hsp90, gp96, and a mixture of two or more of the foregoing.

24. The method according to claim 20 in which the first
5 heat shock protein is hsp70.

25. The method according to claim 20 in which the first
heat shock protein is hsp90.

10 26. The method according to claim 20 in which the first
heat shock protein is gp96.

27. The method according to claim 20 or 21 in which the
first heat shock protein and the second heat shock protein
15 are the same.

28. The method according to claim 20 or 21 in which the
individual is a human.

20 29. The method according to claim 20 in which the first
composition is the same as the second composition.

30. The method according to claim 20 or 21, further
comprising administering to the individual an effective
25 amount of a biological response modifier selected from the
group consisting of interferon- α , interferon- γ , interleukin-
2, interleukin-4, interleukin-6, and tumor necrosis factor.

31. The method according to claim 20 or 21 in which the
30 first composition is administered at weekly intervals.

32. The method according to claim 20 or 21 in which the
second composition is administered at weekly intervals.

35 33. The method according to claim 20 or 21 in which the
first complex is administered intramuscularly,

subcutaneously, intraperitoneally, intravenously, intradermally or mucosally.

34. The method according to claim 20 or 21 in which the
5 second composition is administered intravenously.

35. The method according to claim 20 or 21 in which 10^6
to 10^{12} antigen presenting cells are administered.

10 36. The method according to claim 20, 21 or 23 in which
the first and second complexes are the same.

37. The method according to claim 20 or 21 in which the
cancer comprises a sarcoma or carcinoma, selected from the
15 group consisting of fibrosarcoma, myxosarcoma, liposarcoma,
chondrosarcoma, osteogenic sarcoma, chordoma, angiosarcoma,
endotheliosarcoma, lymphangiosarcoma,
lymphangioendotheliosarcoma, synovioma, mesothelioma, Ewing's
tumor, leiomyosarcoma, rhabdomyosarcoma, colon carcinoma,
20 pancreatic cancer, breast cancer, ovarian cancer, prostate
cancer, squamous cell carcinoma, basal cell carcinoma,
adenocarcinoma, sweat gland carcinoma, sebaceous gland
carcinoma, papillary carcinoma, papillary adenocarcinomas,
cystadenocarcinoma, medullary carcinoma, bronchogenic
25 carcinoma, renal cell carcinoma, hepatoma, bile duct
carcinoma, choriocarcinoma, seminoma, embryonal carcinoma,
Wilms' tumor, cervical cancer, testicular tumor, lung
carcinoma, small cell lung carcinoma, bladder carcinoma,
epithelial carcinoma, glioma, astrocytoma, medulloblastoma,
30 craniopharyngioma, ependymoma, pinealoma, hemangioblastoma,
acoustic neuroma, oligodendroglioma, meningioma, melanoma,
neuroblastoma, retinoblastoma, leukemia, lymphoma, multiple
myeloma, Waldenström's macroglobulinemia, and heavy chain
disease.

35

38. The method according to claim 23 in which the
individual is a human.

39. The method according to claim 20 in which the first antigenic molecule is a peptide with which the first heat shock protein is endogenously associated *in vivo*, and the first complex is prepared from cancerous tissue autologous to the individual.

40. The method according to claim 20 in which the second antigenic molecule is a peptide with which the second heat shock protein is endogenously associated *in vivo*, and the second complex is prepared from cancerous tissue autologous to the individual.

41. The method according to claim 20 in which the first antigenic molecule is a peptide with which the first heat shock protein is endogenously associated *in vivo*, and the first complex is prepared from cancerous tissue allogeneic to the individual.

42. The method according to claim 20 in which the second antigenic molecule is a peptide with which the second heat shock protein is endogenously associated *in vivo*, and the second complex is prepared from cancerous tissue allogeneic to the individual.

43. The method according to claim 20 in which the first antigenic molecule is a peptide with which the first heat shock protein is endogenously associated *in vivo*, and the first complex is prepared from cancerous tissue.

44. The method according to claim 20 in which the second antigenic molecule is a peptide with which the second heat shock protein is endogenously associated *in vivo*, and the second complex is prepared from cancerous tissue.

45. The method according to claim 20 in which the first complex of the first hsp and first antigenic molecule is produced *in vitro*.

46. The method according to claim 20 in which the second complex of the second hsp and second antigenic molecule is produced *in vitro*.

5 47. The method according to claim 20 in which the first antigenic molecule is a tumor-specific antigen.

48. The method according to claim 20 in which the second antigenic molecule is a tumor-specific antigen.

10

49. The method according to claim 20 in which the first complex is purified to 60 to 100 percent of the total mg protein.

15 50. The method according to claim 21 in which the first complex is purified to 60 to 100 percent of the total mg protein.

51. A method of preventing cancer in an individual in
20 whom prevention of cancer is desired comprising:

- (a) administering to the individual a first composition comprising a first complex consisting essentially of a first heat shock protein noncovalently bound to a first antigenic molecule; and
 - 25 (b) administering to the individual a second composition comprising antigen presenting cells sensitized *in vitro* with a sensitizing amount of a second complex of a second heat shock protein noncovalently bound to a second antigenic molecule;
- 30 in which said second composition is administered before, concurrently or after administration of the first composition.

52. The method according to claim 51 in which the first
35 heat shock protein is selected from the group consisting of hsp70, hsp90, gp96, and a mixture of two or more of the foregoing.

53. The method according to claim 51 in which the second heat shock protein is selected from the group consisting of hsp70, hsp90, gp96, and a mixture of two or more of the foregoing.

5

54. The method according to claim 52 in which the second heat shock protein is selected from the group consisting of hsp70, hsp90, gp96, and a mixture of two or more of the foregoing.

10

55. The method according to claim 51 in which the first heat shock protein is hsp70.

56. The method according to claim 51 in which the first
15 heat shock protein is hsp90.

57. The method according to claim 51 in which the first heat shock protein is gp96.

20 58. The method according to claim 51 or 52 in which the first heat shock protein and the second heat shock protein are the same.

59. The method according to claim 51 or 52 in which the
25 individual is a human.

60. The method according to claim 51 in which the first composition is the same as the second composition.

30 61. The method according to claim 51 or 52 in which the first and second complexes are the same.

62. The method according to claim 51 in which the first antigenic molecule is a peptide with which the first heat
35 shock protein is endogenously associated in vivo.

63. The method according to claim 51 in which the second antigenic molecule is a peptide with which the second heat shock protein is endogenously associated *in vivo*.

5 64. The method according to claim 51 in which the first complex is prepared from cancerous tissue.

65. The method according to claim 51 in which the second complex is prepared from cancerous tissue.

10

66. The method according to claim 51 in which the first complex of the first heat shock protein and first antigenic molecule is produced *in vitro*.

15 67. The method according to claim 51 in which the second complex of the second heat shock protein and second antigenic molecule is produced *in vitro*.

68. The method according to claim 51 in which the first
20 antigenic molecule is a tumor-specific antigen.

69. The method according to claim 51 in which the second antigenic molecule is a tumor-specific antigen.

25 70. A method of treating or preventing an infectious disease in an individual in whom such treatment or prevention is desired comprising:

- 30 (a) administering to the individual a first composition comprising a first complex consisting essentially of a first heat shock protein noncovalently bound to a first antigenic molecule; and
- 35 (b) administering to the individual a second composition comprising antigen presenting cells sensitized *in vitro* with a sensitizing amount of a second complex of a second heat shock protein noncovalently bound to a second antigenic molecule;

in which said second composition is administered before, concurrently or after administration of the first composition.

5 71. The method according to claim 70 in which the first heat shock protein is selected from the group consisting of hsp70, hsp90, gp96, and a mixture of two or more of the foregoing.

10 72. The method according to claim 70 in which the second heat shock protein is selected from the group consisting of hsp70, hsp90, gp96, and a mixture of two or more of the foregoing.

15 73. The method according to claim 71 in which the second heat shock protein is selected from the group consisting of hsp70, hsp90, gp96, and a mixture of two or more of the foregoing.

20 74. The method according to claim 70 in which the first heat shock protein is hsp70.

 75. The method according to claim 70 in which the first heat shock protein is hsp90.

25 76. The method according to claim 70 in which the first heat shock protein is gp96.

 77. The method according to claim 70 or 71 in which the
30 first heat shock protein and the second heat shock protein are the same.

 78. The method according to claim 70 or 71 in which the individual is a human.

35

79. The method according to claim 70 in which the first composition is the same as the second composition.

80. The method according to claim 70 or 71 in which the 5 first and second complexes are the same.

81. The method according to claim 70 in which the first antigenic molecule is a peptide with which the first heat shock protein is endogenously associated in cells infected 10 with an infectious agent that causes the infectious disease.

82. The method according to claim 70 in which the second antigenic molecule is a peptide with which the second heat shock protein is endogenously associated in cells 15 infected with an infectious agent that causes the infectious disease.

83. The method according to claim 70 in which the first antigenic molecule is an antigen of an infectious agent that 20 causes the infectious disease.

84. The method according to claim 70 in which the second antigenic molecule is an antigen of an infectious agent that causes the infectious disease. 25

85. The method according to claim 83 or 84 in which the infectious agent is a virus, bacterium, protozoa, fungus, or parasite.

30 86. The method according to claim 1, 20, 51 or 70 in which the antigen presenting cells comprise macrophages.

87. A kit comprising in a first container a composition comprising a complex consisting essentially of a heat shock 35 protein noncovalently bound to an antigenic molecule and a pharmaceutical carrier; and in a second container human antigen presenting cells.

88. The kit of claim 87 in which the heat shock protein is selected from the group consisting of hsp70, hsp90, gp96, and a mixture of two or more of the foregoing.

5 89. The method according to claim 1, 20, 51 or 70 in which the first complex is purified to apparent homogeneity as detected by sodium dodecyl sulfate-polyacrylamide gel electrophoresis.

10 90. The kit according to claim 87 in which the first complex is purified to apparent homogeneity as detected by sodium dodecyl sulfate-polyacrylamide gel electrophoresis.

15

20

25

30

35

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/02194

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : A61K 39/385, 39/39; C07K 1/10

US CL : 424/193.1, 196.11, 197.11; 530/403

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/193.1, 196.11, 197.11; 530/403

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, DIALOG biotech cluster

terms: heat shock, cancer, infectious, immune, response, adjuvant, hsp70, hsp90, gp96

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	GARBE, T.R. Heat shock proteins and infection: Interactions of pathogen and host. Experientia. 15 July 1992, Vol. 48, No. 7, pages 635-639, see entire document.	1-86
A,P ---- X,P	TAMURA, Y. et al. Immunotherapy of tumors with autologous tumor-derived heat shock protein preparations. Science. 3 October 1997, Vol. 278, pages 117-121, see entire document.	1-69, 89 ---- 87, 88, 90
A ---- X	UDONO, H. et al. Cellular requirements for tumor-specific immunity elicited by heat shock proteins: Tumor rejection antigen gp96 primes CD8+ T cells in vivo. Proc. Natl. Acad. Sci. (USA). April 1994, Vol.91, pages 3077-3081, see entire document.	1-69, 89 ---- 87, 88, 90

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
B earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

Date of mailing of the international search report

08 MAY 1998

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Authorized officer

Telephone No.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/02194

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	ARNOLD, D. et al. Cross-priming of minor histocompatibility antigen-specific cytotoxic T cells upon immunization with the heat shock protein gp96. J. Exp. Med. September 1995, Vol. 182, pages 885-889, see entire document.	1-69, 87-90
A	FEIGE, U. et al. Heat shock proteins. Experientia. 15 July 1992, Vol. 48, No. 7, pages 621-622, see entire document.	1-90

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/02194

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☒ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/02194

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains claims directed to more than one species of the generic invention. These species are deemed to lack Unity of Invention because they are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for more than one species to be searched, the appropriate additional search fees must be paid. The species are as follows:

- I. A method for treating or preventing cancer using heat shock proteins.
- II. A method for treating or preventing infectious disease using heat shock proteins.

The claims are deemed to correspond to the species listed above in the following manner:

Species I - claims 20-69.

Species II - claims 70-85.

The following claims are generic: 1-19 and 86-90.

The species listed above do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, the species lack the same or corresponding special technical features for the following reasons: While the species share the same method of practice, the skilled artisan would not reasonably predict that the same method would be effective for use against both cancer and infectious disease. Each condition encompasses a variety of species which have distinct modes of action and elicit distinct immune responses by the affected individual/host.

[Return to PatentWeb top page](#)

[Click Here for Current Status of Order](#)

Order Reference Number: 5721073

2002/04/19 01:42:43 GMT: ORDER SUBMITTED

2002/04/19 01:42:45 GMT: Preparing and emailing patent number US6361521, 15 pages

Delivery mode: Email PDF

Patent number US6361521 [mpo 105779 paedpharm], 15 pages: \$ 4.95

Order Total: \$ 4.95

Connecticut residents will be charged 6% sales tax.

Please Note:

The patents in an email order are sent as individual email messages.

Email orders in plain text or HTML formats are normally processed at a rate of less than thirty seconds per patent. PDF orders may take longer. You may click "Current Status" to see how the preparation of your order is progressing. Remember that a patent that has been emailed to you will generally not be available instantly in your mailbox. The delay depends on the configuration of the relay programs involved in email delivery from MicroPatent® to you.

IMPORTANT NOTE: Corporate email systems typically impose a size quota for individual email messages or the cumulative contents of your mailbox. If the processing of your order was successful, but email delivery failed due to a size problem on your end or an out of date email address we have on file for you (please keep your account information up to date!), the patents you ordered may be downloaded via "Retrieve Orders" on the login page or on the PatentWeb Services page.

The preparation of your order will proceed even if your session expires or is disrupted. There is no need to re-enter your order.

[Return to PatentWeb top page](#)